arXiv:1609.08721v1 [math.KT] 28 Sep 2016

CHOW RINGS OF VERSAL COMPLETE FLAG VARIETIES

NOBUAKI YAGITA

Abstract. In this paper, we compute Chow rings of generically twisted (versal) complete flag varieties corresponding to simple Lie groups.

1. Introduction

Fix a prime number p. Let k be a field of $ch(k) = 0$ and $\bar{k}$ its algebraic closure. Let X be a flag variety over k and $\bar{X} = X|_{\bar{k}}$. We study non-zero elements x such that

$$x \in Ker(res : CH^*(X)/p \to CH^*(\bar{X})/p).$$

Elements in the Chow group of the (original) Rost motive defined by Rost, Voevodsky ([Ro1,2], [Vo2,4]) give such examples. In this paper, we study $CH^*(X)/p$ and give another examples.

Let G and T be a connected compact Lie group and its maximal torus. Let G_k and T_k be a split reductive group and split maximal torus over k corresponding to Lie groups G and T. Let B_k be the Borel subgroup containing T_k.

Moreover we take k such that there is a G_k-torsor $\mathbb{G}_k$ which is isomorphic to a versal G_k-torsor (for the definition, see [Se-Me-Ga], [Me-Ne-Za], [Ka], [To2]), namely, $X = \mathbb{G}_k/B_k$ is the *most twisted* complete flag variety. (We say such X as a generically twisted or a versal flag variety [Me-Ne-Za], [Ka].) In this paper, we study the Chow ring $CH^*(X)$ for the versal flag variety X.

By Petrov-Semenov-Zainoulline ([Pe-Se-Za], [Se-Zh], [Se]), it is known that the p-localized motive $M(X)_{(p)}$ of X is decomposed as

$$M(X)_{(p)} = M(\mathbb{G}_k/B_k)_{(p)} \cong \oplus_i R(\mathbb{G}_k) \otimes \mathbb{T}^{\otimes s_i}$$

where $\mathbb{T}$ is the Tate motive and $R(\mathbb{G}_k)$ is a some motive called generalized Rost motive. (It is the original Rost motive when G is of type

2010 *Mathematics Subject Classification.* 55N20, 55T25, 19L50.

Key words and phrases. Chow ring, Rost motive, complete flag variety.

(I) explained below). Hence we have maps

$$CH^*(BB_k) \to CH^*(X) \to CH^*(R(\mathbb{G}_k))$$

where BB_k is the classifying space for B_k-bundles.

Remark. In this paper, a map $A \to B$ (resp. $A \cong B$) for rings A, B is always a ring map (resp. ring isomorphism). However $CH^*(R(\mathbb{G}_k))$ does not have the canonical ring structure, in general. Hence a map $A \to CH^*(R(\mathbb{G}))/p$ (resp. $A \cong CH^*(R(\mathbb{G}_k))/p$) means only a (graded) additive map (resp. additive isomorphism). For example, the above first map is a ring map but the second is not a ring map.

From Merkurjev and Karpenko [Me-Ne-Za], [Ka], we know that the first map is also surjective. We study what elements in $CH^*(BB_k) \cong CH^*(BT_k)$ generates $CH^*(R(\mathbb{G}_k))$.

Our results are

Theorem 1.1. *For $(G,p) = (SO(2\ell+1), 2)$ and a versal flag variety $X = \mathbb{G}_k/B_k$, there are isomorphisms*

$$CH^*(R(\mathbb{G}_k))/2 \cong \mathbb{Z}/2[c_1, ..., c_\ell]/(c_1^2, ..., c_\ell^2) = \Lambda(c_1, ..., c_\ell),$$

$$CH^*(X)/2 \cong S(t)/(2, c_1^2, ..., c_\ell^2)$$

where $c_i = \sigma_i(t_1, ..., t_\ell)$ is the i-th elementary symmetric function in

$$S(t) = CH^*(BB_k) \cong H^*(BT) \cong \mathbb{Z}[t_1, ..., t_\ell]$$

(i.e., i-th Chern class for the representation $T \subset U(\ell)$).

Remark that we have an isomorphism $CH^*(X)/2 \cong H^*(Sp(\ell)/T; \mathbb{Z}/2)$ for the symplectic group $Sp(\ell)$ (see Corollary 7.7).

Theorem 1.2. *([Ya4]) For G of type (I) and $rank(G) = \ell$, there are isomorphisms*

$$CH^*(R(\mathbb{G}_k))/p \cong \mathbb{Z}/p\{b_1, ..., b_{2p-2}\},$$

$$CH^*(X)/p \cong S(t)/(p, b_i b_j, b_k | 0 \le i, j \le 2p-2 < k \le \ell)$$

where $\mathbb{Z}/p\{a, b, ...\}$ is the $\mathbb{Z}/p$-free module generated by $a, b, ...$ Here $b_i \in H^*(BT)$ are transgression images in the spectral sequence induced from the fibering $G \to G/T \to BT$.

The above b_i are explicitly known ([Tod2], [Tod-Wa], [Na]), for example, when $(G,p) = (G_2, 2)$, we can take $b_1 = t_1^2 + t_1 t_2 + t_2^2$ and $b_2 = t_2^3$ in $H^*(BT) \cong \mathbb{Z}[t_1, t_2]$ with $|t_i| = 2$.

To explain the transgression and type (I) groups, we recall how to compute $H^*(G/T)$ in algebraic topology. By Borel, its mod p-cohomology is (for p odd)

$$H^*(G; \mathbb{Z}/p) \cong P(y)/p \otimes \Lambda(x_1, ..., x_\ell),$$

where $|x_i| = odd$ and $|x_i| \leq |x_j|$ for $i < j$, and where $P(y)$ is a truncated polynomial generated by *even* dimensional elements y_i. When $p = 2$, we consider the graded ring $grH^*(G; \mathbb{Z}/2)$ which is isomorphic to the right hand side ring above.

When G is simply connected and $P(y)$ is generated by just one generator, we call that G is of type (I). Except for $(E_7, p = 2)$ and $(E_8, p = 2, 3)$, all exceptional (simple) Lie groups are of type (I). Note that in these cases, we know $rank(G) = \ell \geq 2p - 2$.

We consider the fibering ([Tod2], [Mi-Ni]) $G \xrightarrow{\pi} G/T \xrightarrow{i} BT$ and the induced spectral sequence

$$E_2^{*,*} = H^*(BT; H^*(G; \mathbb{Z}/p)) \Longrightarrow H^*(G/T; \mathbb{Z}/p).$$

Here we can write $H^*(BT) \cong S(t) = \mathbb{Z}[t_1, ..., t_\ell]$ with $|t_i| = 2$.

It is well known that $y_i \in S(y)$ are permanent cycles and that there is a regular sequence $(\bar{b}_1, ..., \bar{b}_\ell)$ in $H^*(BT)/(p)$ such that $d_{|x_i|+1}(x_i) = \bar{b}_i$ ([Tod2], [Mi-Ni]). The element $\bar{b}_i$ is called the transgression image of x_i. We know that G/T is a manifold of torsion free, and we see that there is a filtration in $H^*(G/T)_{(p)}$ such that

$$grH^*(G/T)_{(p)} \cong P(y) \otimes S(t)/(b_1, ..., b_\ell)$$

where $b_i = \bar{b}_i \ mod(p)$.

The transgression images b_i in Theorem 1.2 are just b_i above. When $(G, p) = (SO(2\ell + 1), 2)$ we can take $b_i = c_i$. In fact, $b_i \ mod(p)$ are generators (such that $|b_i| \leq |b_j|$ for $i < j$) of the kernel $I(p)$ of the map $H^*(BT)/p \to H^*(G/T)/p$ (that is, $CH^*(BB_k)/p \to CH^*(G_k/B_k)/p$).

By giving the filtration on $S(t)$ by b_i, we have the (additive) isomorphism

$$grS(t)/p \cong A \otimes S(t)/(b_1, ..., b_\ell) \quad for\ A = \mathbb{Z}/p[b_1, ..., b_\ell].$$

In particular, we have maps

$$A \xrightarrow{i_A} CH^*(X)/p \to CH^*(R(\mathbb{G}_k))/p.$$

We easily see that $i_A(A) \supset CH^*(R(\mathbb{G}_k))/p$. In particular the above composition map is surjective. Using these, we can prove other results.

Theorem 1.3. *Let $G = G_1 \times G_2$ with $G_1 \cong G_2$ of type (I). Let us write $A = CH^*(R(\mathbb{G}_{1,k}))/p \otimes CH^*(R(\mathbb{G}_{2,k}))/p$ which is isomorphic to*

$$\mathbb{Z}/p\{b_s(1)b_r(2) | 0 \leq s, r \leq 2p - 2\}.$$

Letting $b_k(m) = pr_m^*(b_k)$ for the projection $pr_m : G \to G_m$, $m = 1, 2$, we have surjective maps

$$A \twoheadrightarrow CH^*(R(\mathbb{G}_k))/p \twoheadrightarrow A/(b_{2i-1}(1)b_{2j}(2) - b_{2i}(1)b_{2j-1}(2))$$

where i, j range $1 \leq i, j \leq p-1$.

In particular, $b_{2i-1}(1)b_{2i-1}(2)$ is a product of nonzero torsion elements in $CH^*(X)_{(p)}$, and it is not represented by element in the Chow group of the (original) Rost motive.

Theorem 1.4. *Let $(G,p) = (Spin(2\ell+1), 2)$. Then we have the surjective maps*

$$\Lambda(c_2, ..., c_\ell) \twoheadrightarrow CH^*(R(\mathbb{G}_k))/2 \twoheadrightarrow \mathbb{Z}/2\{1, c_2, ..., c_\ell\}.$$

Theorem 1.5. *Let $(G,p) = (E_7, 2)$, $(E_8, 2)$ or $(E_8, 3)$ so that $\ell = 7$ for E_7 and $\ell = 8$ for E_8. Then we have the surjective map*

$$CH^*(R(\mathbb{G}_k))/p \twoheadrightarrow \mathbb{Z}/p\{1, b_1, ..., b_\ell\}.$$

Of course, there are many other non-zero elements in $CH^*(R(\mathbb{G}_k))/p$. For example, when $(G = E_8, p = 3)$, there are non zero elements b_1b_6, b_1b_8, b_2b_8.

To see the above elements are non zero, we use the torsion index $t(G)_{(p)}$. For $dim(G/T) = 2d$, the torsion index is defined as

$$t(G) = |H^{2d}(G/T; \mathbb{Z})/H^{2d}(BT; \mathbb{Z})|.$$

Let $n(\mathbb{G}_k)$ be the greatest common divisor of the degrees of all finite field extension k' of k such that $\mathbb{G}_k$ becomes trivial over k'. Then by Grothendieck [Gr], it is known that $n(\mathbb{G}_k)$ divides $t(G)$. Moreover, when $\mathbb{G}_k$ is a versal G_k-torsor, we have $n(\mathbb{G}_k) = t(G)$ ([To2], [Ga-Me-Se]). Totaro determined [To1,2]) torsion indexes for all simply connected compact Lie groups G. In particular, $t(E_8) = 2^6 3^2 5$. We also give its another proof.

The plan of this paper is the following. In §2, §3, we recall and prepare the topological arguments for $H^*(G/T)$ and $BP^*(G/T)$. In §4, we recall the decomposition of the motive of a generically twisted flag variety. In §5, we recall the torsion index briefly. In §6, we study $U(m), Sp(m)$ and $PU(p)$ for each p. In §7, §8 we study in the cases $SO(m)$ and $Spin(m)$ for $p = 2$. In §9, §10, we study the cases that G is of type (I) and its product. In §11, §12, §13, we study the cases $(G,p) = (E_8, 3), (E_8, 2)$ and $(E_7, 2)$ respectively.

2. Lie groups G and the flag manifolds G/T

Let G be a connected compact Lie group. By Borel, its mod p-cohomology is (for p odd)

$$H^*(G; \mathbb{Z}/p) \cong P(y)/p \otimes \Lambda(x_1, ..., x_\ell), \quad \ell = rank(G) \tag{2.1}$$
$$with \;\; P(y) = \mathbb{Z}_{(p)}[y_1, ..., y_k]/(y_1^{p^{r_1}}, ..., y_k^{p^{r_k}})$$

where the degree $|y_i|$ of y_i is even and $|x_j|$ is odd. When $p = 2$, a graded ring $grH^*(G;\mathbb{Z}/2)$ is isomorphic to the right hand side ring, e.g. $x_j^2 = y_{i_j}$ for some y_{i_j}. In this paper, $H^*(G;\mathbb{Z}/2)$ means this $grH^*(G;\mathbb{Z}/2)$ so that (2.1) is satisfied also for $p = 2$.

Let T be the maximal torus of G. and BT be the classifying space of T. We consider the fibering ([Tod2], [Mi-Ni])

$$(2.1) \quad G \xrightarrow{\pi} G/T \xrightarrow{i} BT$$

and the induced spectral sequence

$$E_2^{*,*} = H^*(BT; H^*(G;\mathbb{Z}/p)) \Longrightarrow H^*(G/T;\mathbb{Z}/p).$$

The cohomology of the classifying space of the torus is given by

$$H^*(BT) \cong S(t) = \mathbb{Z}[t_1, ..., t_\ell] \quad with \quad |t_i| = 2,$$

where $t_i = pr_i^*(c_1)$ is the 1-st Chern class induced from

$$T = S^1 \times ... \times S^1 \xrightarrow{pr_i} S^1 \subset U(1).$$

for the i-th projection pr_i. Note that $\ell = rank(G)$ is also the number of the odd degree generators x_i in $H^*(G;\mathbb{Z}/p)$.

It is well known that y_i are permanent cycles and that there is a regular sequence ([Tod2],[Mi-Ni]) $(\bar{b}_1, ..., \bar{b}_\ell)$ in $H^*(BT)/(p)$ such that $d_{|x_i|+1}(x_i) = \bar{b}_i$. Thus we get

$$E_\infty^{*,*'} \cong grH^*(G/T;\mathbb{Z}/p) \cong P(y)/p \otimes S(t)/(\bar{b}_1, ..., \bar{b}_\ell).$$

Moreover we know that G/T is a manifold of torsion free, and

$$(2.2) \quad H^*(G/T)_{(p)} \cong \mathbb{Z}_{(p)}[y_1, .., y_k] \otimes S(t)/(f_1, ..., f_k, b_1, ..., b_\ell)$$

where $b_i = \bar{b}_i \ mod(p)$ and $f_i = y_i^{p^{r_i}} \ mod(t_1, ..., t_\ell)$.

Let $BP^*(-)$ be the Brown-Peterson theory with the coefficients ring $BP^* \cong \mathbb{Z}_{(p)}[v_1, v_2, ...]$, $|v_i| = -2(p^i - 1)$ ([Ha], [Ra]). Since $H^*(G/T)$ is torsion free, we also know

$$(2.3) \quad BP^*(G/T) \cong BP^*[y_1, ..., y_k] \otimes S(t)/(\tilde{f}_1, ..., \tilde{f}_k, \tilde{b}_1, ..., \tilde{b}_\ell)$$

where $\tilde{b}_i = b_i \ mod(BP^{<0})$ and $\tilde{f}_i = f_i \ mod(BP^{<0})$.

Let G_k be the split reductive algebraic group corresponding G, and T_k be the split maximal torus corresponding T. Let B_k be the Borel subgroup with $T_k \subset B_k$. Note that G_k/B_k is cellular, and $CH^*(G_k/T_k) \cong CH^*(G_k/B_k)$. Hence we have

$$CH^*(G_k/B_k) \cong H^*(G/T) \quad and \quad CH^*(BB_k) \cong H^*(BT).$$

Let $\Omega^*(-)$ be the BP-version of the algebraic cobordism ([Vo1], [Le-Mo1,2], [Ya2,3])

$$\Omega^*(X) = MGL^{2*,*}(X)_{(p)} \otimes_{MU^*_{(p)}} BP^*$$

$$\text{so that} \quad \Omega^*(X) \otimes_{BP^*} \mathbb{Z}_{(p)} \cong CH^*(X)_{(p)}$$

where $MGL^{*,*'}(X)$ is the algebraic cobordism theory defined by Voevodsky with $MGL^{2*,*}(pt.) \cong MU^*$ the complex cobordism ring. There is a natural (realization) map $\Omega^*(X) \to BP^*(X(\mathbb{C}))$. In particular, we have

$$\Omega^*(G_k/B_k) \cong BP^*(G/T).$$

Let $I_n = (p, v_1, ..., v_{n-1})$ and I_∞ is the (prime invariant) ideal in BP^*. We also note

$$\Omega^*(G_k/B_k)/I_\infty \cong BP^*(G/T)/I_\infty \cong H^*(G/T)/p.$$

3. $BP^*(G/T)$ and $k(n)^*(G/T)$

Recall that $k(n)^*(X)$ is the connected Morava K-theory with the coefficients ring $k(n)^* \cong \mathbb{Z}/p[v_n]$ and $\rho : k(n)^*(X) \to H^*(X; \mathbb{Z}/p)$ is the natural (Thom) map. Recall that there is an exact sequence (Sullivan exact sequence [Ra], [Ya2])

$$... \to k(n)^{*+2(p^n-1)}(X) \xrightarrow{v_n} k(n)^*(X) \xrightarrow{\rho} H^*(X; \mathbb{Z}/p) \xrightarrow{\delta} ...$$

such that $\rho \cdot \delta(x) = Q_n(x)$. Here the Milnor Q_i operation

$$Q_i : H^*(X; \mathbb{Z}/p) \to H^{*+2p^i-1}(X; \mathbb{Z}/p)$$

is defined by $Q_0 = \beta$ and $Q_{i+1} = [P^{p^i} Q_i, Q_i P^{p^i}]$ for the Bockstein β and the reduced powers P^j.

We recall the Serre spectral sequence

$$E_2^{*,*'} \cong H^*(BT; H^*(G; \mathbb{Z}/p)) \Longrightarrow H^*(G/T; \mathbb{Z}/p).$$

induced from the fibering $G \xrightarrow{\pi} G/T \xrightarrow{i} BT$ in the preceding section.

Lemma 3.1. *(Lemma 4.3 in [Ya1]) In the spectral sequence $E_r^{*,*'}$ above, suppose that there is $x \in H^*(G; \mathbb{Z}/p)$ such that*

$$(*) \quad y = Q_n(x) \neq 0 \quad \text{and} \quad b = d_{|x|+1}(x) \neq 0 \in E^{*,0}_{|x|+1}.$$

Moreover suppose that $E^{0,|x|}_{|x|+1} \cong \mathbb{Z}/p\{x\} \cong \mathbb{Z}/p$. Then there are $y' \in k(n)^*(G/T)$ and $b' \in k(n)^*(BT)$ such that $\pi^*(y') = y$, $\rho(b') = b$ and that

$$(**) \quad v_n y' = \lambda i^* b' \in k(n)^*(p^{-1}BT^{|b|}), \quad \lambda \neq 0 \in \mathbb{Z}/p.$$

Conversely if $(**)$ holds in $k(n)^*(G/T)$ for $y = \pi^*(y') \neq 0$ and $b = \rho(b') \neq 0$, then there is $x \in H^*(G; \mathbb{Z}/p)$ such that $(*)$ holds.

Proof. Let $B' = BT^{|b|-1}$ be the $|b|-1$ dimensional skeleton of BT, and $E' = i^{-1}(B')$ where we take i an projection. Consider the Serre spectral sequence

$$H^*(B'; H^*(G; \mathbb{Z}/p)) \Longrightarrow H^*(E'; \mathbb{Z}/p).$$

Since $d_r(x) = b = 0 \in H^*(B'; \mathbb{Z}/p)$, there is $x' \in H^*(E'; \mathbb{Z}/p)$ such that $\pi^*(x') = x$. Let $Q_n(x') = y'$ so that $\pi^* y' = y$. Then by the Sullivan exact sequence $v_n y' = 0$ in $k(n)^*(E')$.

On the other hand, let $B'' = B^{|b|-1} \cup e_b$ and $E'' = i^{-1}B''$ where e_b is the normal cell representing b. Then $d_r x = b \neq 0 \in H^*(B''; \mathbb{Z}/p)$. By the supposition in this lemma, there does not exist $x'' \in H^*(E''; \mathbb{Z}/p)$ such that $i^*(Q_n x'') = y$, that is, for each $y'' \in H^*(E''; \mathbb{Z}/p)$ with $\pi^* y'' = y$, we see $v_n y'' \neq 0 \in k(n)^*(E'')$.

For $j : E' \subset E''$, we can take an element y'' with $j^*(y'') = y'$ by the following reason. Suppose that $\bar{y} = y' - j^* y'' \neq 0 \in H^*(E'; \mathbb{Z}/p)$ is not in $j^* H^*(E''; \mathbb{Z}/p)$. It means $d_r(\bar{y}) = sb \neq 0$ for some $s \in H^*(G; \mathbb{Z}/p)$ and $r < |b|$. However all differential in $E_s^{*,*'}$ are written by $d_{|b_i|}(x_i) = b_i$, and $(b_1, ..., b_\ell)$ is regular in $S(t)$. Hence there is not such $\bar{y}$.

Since $v_n y' = v_n j^*(y'') = 0 \in k(n)^*(E')$ but $v_n y'' \neq 0 \in k(n)^*(E'')$, by dimensional reason, $v_n y'' = \lambda b$ for $\lambda \neq 0 \in \mathbb{Z}/p$.

Conversely, suppose that $v_n y' = i^*(b') \neq 0$ in $k(n)^*(E)$. Then $v_n y' = 0$ in $k(n)^*(E')$ and there is $\tilde{x} \in H^*(E'; \mathbb{Z}/p)$ with $Q_n \tilde{x} = y'$. Then for $\pi^*(y') = y$ and $\pi^*(\tilde{x}) = x$, we see $Q_n(x) = y$. But $\tilde{x}$ does not exist in $H^*(E''; \mathbb{Z}/p)$. Hence $d_{|x|+1}(x) = \lambda b$ for $\lambda \neq 0 \in \mathbb{Z}/p$, by dimensional reason. □

Remark. (Remark 4.8 in [Ya1]) The above theorem also holds letting $k(0)^*(X) = H^*(X; \mathbb{Z})$ and $v_0 = p$. This fact is well known (Lemma 2.1 in [Tod2]).

Corollary 3.2. *Let $b \neq 0$ be the transgression image of x, i.e. $d_{|x|+1}(x) = b$. Then we have the relation*

$$b = \sum_{i=0} v_i y(i) \quad in\ BP^*(G/T)/I_\infty^2$$

where $y(i) \in H^*(G/T; \mathbb{Z}/p)$ with $\pi^* y(i) = Q_i x$.

Proof. Let us write

$$b = py(0) + v_1 y(1) + ... \quad in\ BP^*(G/T)/I_\infty^2.$$

Then $b = v_n y(n)\ mod(v_n^2) \in k(n)^*(G/T)$ where $y(n) \in k(n)^*(G/T)$. However $y(n)$ is decided only with $mod(v_n)$ and we can identify $y(n) \in H^*(G/T; \mathbb{Z}/p)$ by the Sullivan exact sequence. Then we can see the corollary from the above lemma. □

Let G be a *simply* connected Lie group such that $H^*(G)$ has p-torsion. Then it is known ([Mi-Tod]) that $H^*(G)$ has just (not higher) p-torsion in $H^*(G)_{(p)}$. It is also known that there is $m \geq 1$ with

$$(*) \quad P^{p^i}(y_i) = y_{i+1} \quad for\ 1 \leq i \leq m-1, \quad and \quad P^{p^m}(y_m) = 0.$$

(Here suffix i is changed adequately from that defined in the preceding section (2.1). Note $m = 1$ for type (I) groups.) Moreover $|x_1| = 3$ and $P^1(x_1) = -x_2$, and $\beta(x_2) = y_1$. Hence

$$(**) \quad Q_i(x_1) = y_i, \quad Q_0(x_{i+1}) = y_i.$$

Therefore from the preceding corollary, we have

$$v_1 y(1) + ... + v_m y(m) = 0 \quad in\ BP^*(G/T)/I_\infty^2.$$

We will study the above equation more detailedly.

Here we recall the Quillen (Landweber-Novikov) operation ([Ha], [Ra]). For a sequence $\alpha = (a_1, a_2, ...)$, $a_i \geq 0$ with $|\alpha| = \sum_i 2(p^i - 1)a_i$, we have the Quillen operation $r_\alpha : BP^*(X) \to BP^{*+|\alpha|}(X)$ such that

$$(1) \quad \rho(r_\alpha(x)) = \chi P^\alpha(\rho(x)) \quad for\ \rho : BP^*(X) \to H^*(X; \mathbb{Z}/p),$$

where χ is the anti-automorphism in the Steenrod algebra,

$$(2) \qquad r_\alpha(xy) = \sum_{\alpha = \alpha' + \alpha''} r_{\alpha'}(x) r_{\alpha''}(y) \qquad Cartan\ formula,$$

$$(3) \qquad r_{p^i \Delta_{n-i}}(v_n) = \begin{cases} v_i & for\ \Delta_i = (0, ..., 0, \overset{i}{1}, 0, ..., 0). \\ 0 \quad mod(I_\infty^2) & otherwise. \end{cases}$$

We also note that $\Omega^*(X)$ has the same operation r_α satisfying (2),(3) and (1) for $\rho : \Omega^*(X) \to CH^*(X)/p$ and the reduce power operation P^α on $CH^*(X)/p = H^{2*,*}(X; \mathbb{Z}/p)$ defined by Voevodsky.

Lemma 3.3. *If $|\alpha| < |v_{i-1} - v_i| = 2(p^i - p^{i-1})$, then r_α acts on $BP^*(X)/(I_\infty^2, v_i, v_{i+1}, ...)$.*

Proof. When degree has above condition we have $r_\alpha(v_s) \in I_\infty^2$ for all $s \geq i$. □

Let $h^*(-)$ be a cohomology theory (e.g. $H^*(-; \mathbb{Z}/p)$, $k(n)^*(-)$). The product $G \times G \to G$ induces the map

$$\mu : G \times G/T \to G/T.$$

Here note $h^*(G \times G/T) \cong h^*(G) \otimes_{h^*} h^*(G/T)$, since $h^*(G/T)$ is h^*-free. For $x \in h^*(G/T)$, we say that x is *primitive* ([Mi-Ni], [Mi-Tod]) if

$$\mu^*(x) = \pi^*(x) \otimes 1 \ + \ 1 \otimes x \quad where\ \pi : G \to G/T.$$

It is immediate that x is primitive implies so is $r_\alpha(x)$. Of course $b \in BP^*(BT)$ are primitive but by_i are not, in general. We can take y_1 as primitive (adding elements if necessary) in $BP^*(G/T)$.

Lemma 3.4. *Let G be a simply connected Lie group satisfying $(*)$. Let y_1 be a primitive element and define $y_{i+1} = r_{p^i\Delta_1}(y_i)$. Then we have*

$$v_1y_1 + v_2y_2 + ... + v_my_m = b_1 \quad mod(I_\infty^2).$$

Proof. Note that $v_ny(n) = b_1$ is of course primitive in $k(n)^*(G/T)$. We prove $y_n = y(n)\ mod(I_\infty^2)$. Let us write

$$y(n) = y_n + \sum yt \quad with\ y \in P(y),\ t \in S(t),\ |t| \geq 2.$$

We will prove $t = 0$.

Consider the Atiyah-Hirzebruch spectral sequence

$$E_2^{*,*'} \cong H^*(G; k(n)^*) \Longrightarrow k(n)^*(G).$$

The first non-zero differential is $d_{2p^n-1}(x) = v_nQ_n(x)$. Since $|y| \leq |y_n| - 2 = 2p^n$, we see that y is v_n-torsion free in $k(n)^*(G)$. This means if $t \neq 0$, then

$$v_ny \otimes t \neq 0 \quad in\ k(n)^*(G) \otimes k(n)^*(G/T).$$

Therefore $t = 0$ from the primitivity of y_n and $v_ny(n)$. □

Act r_{Δ_1} on the above equation, we have

Lemma 3.5. *In $BP^*(G/T)/(I_\infty^2)$, we have*

$$py_1 + v_1P^1(y_1) + v_2P^1(y_2) + ... + v_mP^1(y_m) = P^1(b_1) = b_2.$$

4. GENERICALLY TWISTED VARIETY

Recall that $\mathbb{G}_k$ is a nontrivial G_k-torsor. We can construct a twisted form of G_k/B_k by

$$(\mathbb{G}_k \times G_k/B_k)/G_k \cong \mathbb{G}_k/B_k.$$

We will study the twisted flag variety $X = \mathbb{G}_k/B_k$.

Let $P \supset T$ be a parabolic subgroup of G. Petrov, Semenov and Zainoulline developed the theory of decompositions of motives $M(\mathbb{G}_k/P_k)$. They develop the theory of generically splitting varieties. We say that L is splitting field of a variety of X if $M(X|_L)$ is isomorphic to a direct sum of twisted Tate motives $\mathbb{T}^{\otimes i}$ and the restriction map $i_L : M(X) \to M(X|_L)$ is isomorphic after tensoring $\mathbb{Q}$. A smooth scheme X is said to be generically split over k if its function field $L = k(X)$ is a splitting field. Note that (the complete flag) $X = \mathbb{G}_k/B_k$ is always generically split, i.e., $X|L$ is cellular.

Theorem 4.1. *(Theorem 3.7 in [Pe-Se-Za]) Let $Q_k \subset P_k$ be parabolic subgroups of G_k which are generically split over k. Then there is a decomposition of motive : $M(\mathbb{G}/Q_k) \cong M(\mathbb{G}_k/P_k) \otimes H^*(P/Q)$.*

By extending the arguments by Vishik [Vi] for quadrics to that for flag varieties, Petrov, Semenov and Zainoulline define the J-invariant of $\mathbb{G}_k$. Recall the expression in §2

$$(*) \quad H^*(G;\mathbb{Z}/p) \cong \mathbb{Z}/p[y_1,...,y_s]/(y_1^{p^{r_1}},...,y_s^{p^{r_s}}) \otimes \Lambda(x_1,...,x_\ell).$$

The J-invariant is defined as $J_p(\mathbb{G}_k) = (j_1,...,j_s)$ if j_i is the minimal integer such that

$$y_i^{p^{j_i}} \in Im(res_{CH}) \quad mod(y_1,...,y_{i-1},t_1,...,t_\ell)$$

for $res_{CH} : CH^*(\mathbb{G}_k/B_k) \to CH^*(G_k/B_k)$. Here we take $|y_1| \le |y_2| \le ...$ in $(*)$. Hence $0 \le j_i \le r_i$ and $J_p(\mathbb{G}_k) = (0,...,0)$ if and only if $\mathbb{G}_k$ split by an extension of the index coprime to p. One of the main results in [Pe-Se-Za] is

Theorem 4.2. *(Theorem 5.13 in [Pe-Se-Za] and Theorem 4.3 in [Se-Zh]) Let $\mathbb{G}_k$ be a G_k-torsor over k, $X = \mathbb{G}_k/B_k$ and $J_p(\mathbb{G}_k) = (j_1,...,j_s)$. Then there is a p-localized motive $R(\mathbb{G}_k)$ such that*

$$M(X) \cong \oplus_u R(\mathbb{G}_k) \otimes \mathbb{T}^{\otimes u}.$$

Here $\mathbb{T}^{\otimes u}$ are Tate motives with $CH^*(\oplus_u \mathbb{T}^{\otimes u})/p \cong P' \otimes S(t)/(b)$ where

$$P' = \mathbb{Z}/p[y_1^{p^{j_1}},...,y_s^{p^{j_s}}]/(y_1^{p^{r_1}},...,y_s^{p^{r_s}}) \subset P(y)/p,$$
$$S(t)/(b) = S(t)/(b_1,...,b_\ell).$$

The $mod(p)$ Chow group of $\bar{R}(\mathbb{G}_k) = R(\mathbb{G}_k) \otimes \bar{k}$ is given by

$$CH^*(\bar{R}(\mathbb{G}_k))/p \cong \mathbb{Z}/p[y_1,...,y_s]/(y_1^{p^{j_1}},...,y_s^{p^{j_s}}).$$

Hence we have $CH^*(\bar{X})/p \cong CH^*(\bar{R}(\mathbb{G}_k)) \otimes P' \otimes S(t)/(b)$ and $CH^*(X)/p \cong CH^*(R(\mathbb{G}_k)) \otimes P' \otimes S(t)/(b)$.

Let P_k be a special (namely, any extension is split, e.g. B_k). Let us consider an embedding of G_k into the general linear group GL_N for some N. This makes GL_N a G_k-torsor over the quotient variety $S = GL_N/G_k$. We define F to be the function field $k(S)$ and define the *versal G_k-torsor* E to be the G_k-torsor over F given by the generic fiber of $GL_N \to S$. (For details, see [Ga-Me-Se], [Me-Ne-Za],[Ka],[To2].)

$$\begin{array}{ccc} E & \longrightarrow & GL_N \\ \downarrow & & \downarrow \\ Spec(k(S)) & \longrightarrow & S = GL_N/G_k \end{array}$$

The corresponding flag variety E/P_k is called *generically twisted* or *versal* flag variety, which is considered as the most complicated twisted flag variety (for given G_k, P_k). It is known that the Chow ring $CH^*(E/P_k)$ is not dependent to the choice of generic G_k-torsors E (Remark 2.3 in [Ka]).

Karpenko and Merkurjev proved the following result (for $CH^*(X)$) for a versal (generically twisted) flag variety.

Theorem 4.3. *(Karpenko Lemma 2.1 in [Ka]) Let $h^*(X)$ be an oriented cohomology theory (e.g., $CH^*(X)$, $\Omega^*(X)$). Let P_k be a parabolic subgroup of G_k and $\mathbb{G}_k/P_k$ be a versal flag variety. Then the natural map $h^*(BP_k) \to h^*(\mathbb{G}_k/P_k)$ is surjective.*

Corollary 4.4. *The Chow ring $CH^*(\mathbb{G}_k/B_k)$ is generated by elements t_i in $S(t)$. In particular, each $x \in CH^*(G_k/B_k)$, the element $p^t x$ is represented by elements in $S(t)$ for $t \geq t(G)_{(p)}$.*

Proof. For some extension F/k, of order ap^t, a is coprime p, the G_k-torsor $\mathbb{G}_k$ splits. Hence $p^t y^i \in Im(res_{CH} : CH^*(\mathbb{G}_k/B_k) \to CH^*(G_k/B_k))$, which is written by elements in $S(t)$ by the above Karpenko theorem □

Corollary 4.5. *If $X = \mathbb{G}_k/B_k$ is a versal flag variety, then $J(\mathbb{G}_k) = (r_1, ..., r_s)$, i.e. $r_i = j_i$.*

Proof. If $j_i < r_i$, then $0 \neq y_i^{p^{j_i}} \in res(CH^*(X) \to CH^*(G_k/T_k))$, which is in the image from $S(t)$ by the preceding theorem. This contradicts to $CH^*(G_k/T_k; \mathbb{Z}/p) \cong P(y)/p \otimes S(t)/(b)$ and $0 \neq y_i^{p^{j_i}} \in P(y)/p$. □

Here we recall the (original) Rost motive R_n defined from a norm variety isomorphic to v_n and nonzero pure symbol in $K^M_{n+1}(k)/p$. When $J(\mathbb{G}_k) = (1)$, we know $R(\mathbb{G}_k) \cong R_2$ from also [Pe-Se-Za]. We write $\bar{R}_n = R_n \otimes \bar{k}$. The Rost motive R_n is defined as a non-split motive but split over a field of degree ap with $(a, p) = 1$, and

$$CH^*(\bar{R}_n) \cong \mathbb{Z}[y]/(y^p), \quad \Omega^*(\bar{R}_n) \cong BP^*[y]/(y^p), \quad |y| = 2(p^n-1)/(p-1).$$

Theorem 4.6. *([Vi-Ya], [Ya3], [Me-Su]) Let R_n be the (original) Rost motive defined by Rost and Voevodsky ([Ro1,2],[Vo2,3]). Then the restriction $res_\Omega : \Omega^*(R_n) \to \Omega^*(\bar{R}_n)$ is injective. Here the restriction image $Im(res_\Omega)$ is isomorphic to*

$$BP^*\{1, v_j y^i \mid 0 \leq j \leq n-1, 1 \leq i \leq p-1\} \subset BP^*[y]/(y^p).$$

Hence writing $v_j y^i = c_j(y^i)$, we have

$$CH^*(R_n)/p \cong \mathbb{Z}/p\{1, c_j(y^i) \mid 0 \leq j \leq n-1,\ 1 \leq i \leq p-1\}.$$

In particular, we have isomorphisms

$$CH^*(R_1)/p \cong \mathbb{Z}/p\{1, py, py^2, ..., py^{p-1}\},$$
$$CH^*(R_2)/p \cong \mathbb{Z}/p\{1, v_1y, py, v_1y^2, py^2, ..., v_1y^{p-1}, py^{p-1}\}.$$

5. TORSION INDEX

Let $dim(G/T) = 2d$. Then the torsion index is defined as

$$t(G) = |H^{2d}(G/T; \mathbb{Z})/H^{2d}(BT; \mathbb{Z})|.$$

Let $n(\mathbb{G}_k)$ be the greatest common divisor of the degrees of all finite field extension k' of k such that $\mathbb{G}_k$ becomes trivial over k'. Then by Grothendieck [Gr], it is known that $n(\mathbb{G}_k)$ divides $t(G)$. Moreover, there is a G_k-torsor $\mathbb{G}_F$ over some extension field F of k such that $n(\mathbb{G}_F) = t(G)$ (in fact, this holds for each versal G_k-torsor [To2], [Me-Ne-Za], [Ka]). Note that $t(G_1 \times G_2) = t(G_1) \cdot t(G_2)$. It is well known that if $H^*(G)$ has a p-torsion, then p divides the torsion index $t(G)$. Torsion index for simply connected compact Lie groups are completely determined by Totaro [To1,2]. For example, $t(E_8) = 2^6 3^2 5$.

Let $\mathbb{G}_k$ be a versal G_k-torsor and $X = \mathbb{G}_k./B_k$ be the versal flag variety. Recall that

$$grH^*(G/T; \mathbb{Z}/p) \cong P(y)/p \otimes S(t)/(b)$$

where $S(t)/(b) = S(t)/(b_1, ..., b_\ell)$, $P(y)/p \cong \mathbb{Z}/p[y_1, ..., y_s]/(y_1^{p^{r_1}}, ..., y_s^{p^{r_s}})$.
Recall Corollary 4.5 and we see $J(\mathbb{G}_k) = (r_1, ..., r_s)$, i.e., $y_i^{p^{r_i}-1} \notin S(t)$.

Giving the filtration on $S(t)$ by b_i, we have the isomorphism

$$grS(t)/p \cong \mathbb{Z}/p[b_1, ..., b_\ell] \otimes S(t)/(b_1, ..., b_\ell).$$

Let us write for $N > 0$

$$A_N = \mathbb{Z}/p\{b_{i_1}...b_{i_k} |\ |b_{i_1}| + ... + |b_{i_k}| \le N\} \subset grS(t).$$

Of course $H^*(G/T) = 0$ for $* > 2d = dim(G/T)$, we have a map

$$S(t)/p \to A_{2d} \otimes S(t)/(b) \to CH^*(X)/p.$$

Lemma 5.1. *The composition map is a surjection*

$$A_{2d} \to CH^*(X)/p \twoheadrightarrow CH^*(R(\mathbb{G}_k))/p.$$

Proof. Recall the decomposition $M(X) \cong \oplus_i R(\mathbb{G}_k) \otimes \mathbb{T}^{s_i}$. Since the restriction map $res_{CH}(CH^*(\mathbb{T}^{s_i})/p \to CH^*(\bar{\mathbb{T}}^{s_i})/p)$ is isomorphic, we have

$$CH^*(\oplus_i \mathbb{T}^{s_i})/p \cong CH^*(\oplus_i \bar{\mathbb{T}}^{s_i})/p$$
$$\cong CH^*(G_k/T_k)/(p, P(y)^+) \cong S(t)/(p, b).$$

Thus we can write $CH^*(\mathbb{T}^{s_i}) \cong \mathbb{Z}_{(p)}\{u_i\}$ for some $u_i \neq 0 \in S(t)/(p,b)$. Hence $CH^*(X)/p$ is generated by elements the product $b{\cdot}u$ in $CH^*(X)/p$ for $b \in CH^*(R(\mathbb{G}_k)) \subset CH^*(X)/p$ and $u \in S(t)/(p,b)$. Note $bu \neq 0$ if $b \neq 0$.

On the other hand, since $CH^*(X)$ is generated by images from $S(t)$, which is generated by $b'u$ for $b' \in Im(A_d \to CH^*(X)/p)$. When $s_i \neq 0$ (i.e., $|u| \geq 2$), then $pr(bu) = 0$ for the projection $CH^*(X)/p \to CH^*(\mathbb{G}_k))/p$. Hence we have the lemma. □

From the arguments in the preceding lemma, we have

Corollary 5.2. *Let $b \in Ker(pr : CH^*(X)/p \to CH^*(R(\mathbb{G}_k))/p)$. Then we can write*

$$b = \sum b'u' \quad with\ b' \in A_{2d},\ u' \in S(t), \quad |u'| > 0.$$

Corollary 5.3. *If $b_i \neq 0$ in $CH^*(X)/p$, then so in $CH^*(R(\mathbb{G}_k))/p$.*

Proof. If $b_i = \sum b'u'$, then $b' \in Ideal(b_1, ..., b_{i-1})$ by $|u'| > 0$. This contradict to that $(b_1, ..., b_\ell)$ is regular. □

Let us write

$$y_{top} = \Pi_{i=1}^{s} y_i^{p^{r_i}-1} \quad (reps.\ t_{top})$$

the generator of the highest degree in $P(y)$ (resp. $S(t)/(b)$) so that $f = y_{top}t_{top}$ is the fundamental class in $H^{2d}(G/T)$.

Lemma 5.4. *The following map is surjective*

$$A_N \twoheadrightarrow CH^*(R(\mathbb{G}_k))/p \quad where\ N = |y_{top}|.$$

Proof. In the preceding lemma, $A_N \otimes u$ for $|u| > 0$ maps zero in $CH^*(R(\mathbb{G}_k))/p$. Since each element in $S(t)$ is written by an element in $A_N \otimes S(t)/(b)$, we have the corollary. □

The following lemma is only used for $G = SO(m)$, which is not simply connected.

Lemma 5.5. *In $H^*(G/T)$, let $b_I = b_{i_1}...b_{i_k}$ and $y(I) = y(i_1)...y(i_k)$ with $py(i_s) = b_{i_s}$. Moreover for all J such that $|b_J| \leq |b_I|$, we assume $y(J) \neq 0 \in P(y)/p$ and $b_I \neq 0 \in CH^*(X)/p$. Then we have $b_I \neq 0 \in CH^*(R(\mathbb{G}_k)/p$.*

Proof. Suppose $b_I \in Ker(pr)$ and let us write

$$b_I = \sum b_J u', \quad b_J \in A_{2d},\ u' \in S(t)/(b), \quad |u'| > 0.$$

Then we can write in $H^*(G/T)$

$$p^k y(I) = \sum p^{s(J,u')} y(J)u' \quad for\ y(J) \notin P(y)/p,\ s(J,u') \geq 1.$$

Let $s = min(k, s(J, u'))$. Since $H^*(G/T)$ has no p-torsion, we have

$$p^{k-s}y(I) = \sum p^{s(J,u')-s}y'u'.$$

When $k = s$, this induces that there is a relation

$$y(I) - \sum p^{s(J,u')-k}y(J)u' = 0 \ \ in \ \ H^*(G/T)/p, \ with \ |u'| > 0.$$

This contradicts to $H^*(G/T)/p \cong P(y)/p \otimes S(t)/(b)$. When $s = s(J', u'')$, there is a relation $y(J')u'' + ... = 0$ in $H^*(G/T)/p$ and it is also a contradiction. □

Now we consider the torsion index.

Lemma 5.6. *Let $\tilde{b} = b_{i_1}...b_{i_k}$ in $S(t)$ such that in $H^*(G/T)$*

$$\tilde{b} = p^s(y_{top} + \sum yt), \quad |t| > 0$$

for some $y \in P(y)$ and $t \in S(t)$. Then $t(G)_{(p)} \leq p^s$.

Proof. Suppose $p^s < t(G)_{(p)}$. We can assume $t(G) = p^{s+1}$ multiplying p^i if necessary. Since $tt_{top} = 0 \in S(t)/(b)$, we see

$$tt_{top} \in Ideal(b_1, ..., b_\ell) \subset Ideal(p).$$

Therefore $p^s \sum ytt_{top} \in Ideal(p^{s+1})$. So it is in $S(t)$, by Karpenko's theorem. So $t(G) \leq p^s$ and this is a contradiction, □

Corollary 5.7. *In the preceding lemma, assume $p^s = t(G)_{(p)}$. Then for each subset $(i'_1, ..., i'_{k'}) \subset (i_1, ..., i_k)$, the element $b'_{i'_1}...b'_{i_{k'}} \neq 0 \in CH^*(X)/p$.*

Proof. Let us write $I' = (i'_1, ..., i'_{k'}) \subset I = (i_1, ..., i_k)$, $I' \cup I'' = I$, and $b_I = b_{i_1}...b_{i_k}$. It is immediate $b_{I'} \notin CH^*(X)/p$ since $b_I = b_{I'}b_{I''} \neq 0 \in CH^*(X)/p$. □

So when $t(G)_{(p)}$ is big enough and there is $\tilde{b}$ in the preceding lemma, we can find many non zero elements in $CH^*(X)/p$ whose restriction images are zero in $CH^*(\bar{X})/p$.

6. The groups $GL(n)$, $Sp(n)$ and $PU(p)$.

We consider the Lie group $G = U(\ell)$, while its cohomology has no torsion. Recall that

$$H^*(U(\ell)) \cong \Lambda(x_1, ..., x_\ell) \quad with \ |x_i| = 2i - 1.$$

So $P(y)/p \cong \mathbb{Z}/p$, and $CH^*(R(\mathbb{G}_k)/p \cong CH^*(\bar{R}(\mathbb{G}_k))/p \cong \mathbb{Z}/p$, that is, there is no twisted form of G_k/B_k. Moreover

$$CH^*(X)/p \cong S(t)/(p, b_1, ..., b_\ell)$$

for $d_{|x_i|+1}(x_i) = b_i$. It is well known that we can take $b_i = c_i$ the i-th elementary symmetric function on $S(t) \cong \mathbb{Z}[t_1, ..., t_\ell]$

Proposition 6.1. *Let $G = GL_\ell$ and p is a prime number. Let $X = G_k/B_k$. Then*

$$CH^*(X)/p \cong S(t)/(p, c_1, ..., c_\ell)$$

where c_i is the Chern class in $H^*(BT) \cong S(t)$ by the map $T \subset U(\ell)$.

Proof. We consider the fibering $G/T \to BT \to BG$. The composition of the induced map

$$H^*(BG) \to H^*(BT) \to H^*(G/T)$$

is zero. The first map induces the isomorphism

$$H^*(BG) \cong H^*(BT)^{W_G(T)} \cong \mathbb{Z}[c_1, ..., c_\ell]$$

Thus $(b_1, ..., b_\ell) \supset (c_1, ..., c_\ell)$. By dimensional reason, we have the proposition. □

Next consider in the case $G' = Sp(\ell)$ and Recall that

$$H^*(Sp(\ell)) \cong \Lambda(x'_1, ..., x'_\ell) \quad with\ |x_i| = 4i - 1.$$

So $P(y)'/p \cong \mathbb{Z}/p$, and there is no twisted form of G'_k/B_k. Moreover we have $d_{|x'_i|+1i}(x'_i) = p_i$ the Pontryagin class. Hence we have

Proposition 6.2. *Let $G' = Sp(\ell)$ and p is a prime number. Let $X' = G'_k/B_k$. Then*

$$CH^*(X')/p \cong S(t)/(p, p_1, ..., p_\ell).$$

In particular, we have $CH^*(X')/2 \cong S(t)/(2, c_1^2, ..., c_\ell^2)$.

Now we consider in the case $(G, p) = (PU(p), p)$. Its mod p cohomology is

$$H^*(G; \mathbb{Z}/p) \cong \mathbb{Z}/p[y]/(y^p) \otimes \Lambda(x_1, ..., x_{p-1}) \quad |y| = 2,\ |x_i| = 2i - 1.$$

So $P(y)/p \cong \mathbb{Z}/p[y]/(y)$ with $|y| = 2$. Since G is not simply connected, G is not of type I while $P(y)$ is generated by only one y. (However $CH^*(X)/p$ is quite resemble to that of type (I). Compare Theoem 6.4 and Theorem 9.4 below.)

We consider the map $U(p-1) \to U(p) \to PU(p)$ where the maximal tori of $U(p-1)$ and $PU(p)$ are isomorphic, i.e., $T_{U(p-1)} \cong T_{PU(p)}$. By using the map $U(p-1) \to PU(p)$, we know $d_{2i}(x_i) = c_i$. Hence we have

$$grH^*(G/T; \mathbb{Z}/p) \cong \mathbb{Z}/p[y]/(y^p) \otimes S(t)/(c_1, ..., c_{p-1}).$$

Lemma 6.3. *We have $py^i = c_i \in H^*(G/T)_{(p)}$, and hence $\beta(x_i) = y^i \in H^*(G; \mathbb{Z}/p)$.*

Proof. Since $H^*(G/T)$ is torsion free, if $py^i \notin S(t)$, then we can inductively write

$$py^i = \sum_{0<s<i} y^s t(s) \quad for\ some\ s,\ t(s) \neq 0 \in S(t)/(p, c_1, ..., c_{i-1}).$$

But it contradicts to $H^*(G/T)/p \cong P(y) \otimes S(t)/(p, b)$ where $b_i = c_i$. □

Theorem 6.4. *Let $G = PU(p)$ and $X = \mathbb{G}_k/B_k$. Then there are isomorphisms*

$$CH^*(R(\mathbb{G}_k))/p \cong CH^*(R_1)/p \cong \mathbb{Z}/p\{1, c_1, ..., c_{p-1}\},$$

$$CH^*(X)/p \cong S(t)/(p, c_i c_j | 1 \leq i, j \leq p-1).$$

Proof. It is known from [Pe-Se-Za] that $R(\mathbb{G}_k) \cong R_1$. Hence the second isomorphism follows from $py^i = c_i$ and Theorem 4.6

$$CH^*(R_1)/p \cong \mathbb{Z}/p\{1, py, ..., py^{p-1}\}.$$

From the main theorem of [Pe-Se-Za], we have the additive isomorphism

$$CH^*(X)/p \cong \mathbb{Z}/p\{1, py, ..., py^{p-1}\} \otimes S(t)/(b) \quad where\ b_i = c_i.$$

Note that $c_i c_j = p^2 y^{i+j} = p c_{i+j}$ in $\Omega^*(\bar{X})$. Since $res_\Omega : \Omega^*(X) \to \Omega^*(\bar{X})$ is injective, we see $c_i c_j = 0 \in CH^*(X)/p$.

Of course we have an additive isomorphism

$$S(t)/(p, c_i c_j) \cong \mathbb{Z}/p\{1, c_1, ..., c_{p-1}\} \otimes S(t)/(c_1, ..., c_{p-1}).$$

Moreover we have a surjective ring map $S(t)/(p, c_i c_j) \twoheadrightarrow CH^*(X)/p$. From the additive isomorphism, its kernel is zero, which induces the ring isomorphism of the theorem. □

We note here the following lemma for a (general) split algebraic group G_k and a G_k-torsor $\mathbb{G}_k$.

Lemma 6.5. *The composition of the following map*

$$CH^*(BG_k)/p \to CH^*(BB_k)/p \to CH^*(\mathbb{G}_k/B_k)/p$$

is zero for $* > 0$.

Proof. Take U (e.g., GL_N for a large N) such that U/G_k approximates the classifying space BG_k. Namely, we can take $\mathbb{G}_k = f^*U$ for the

classifying map $f : \mathbb{G}_k/G_k \to U/G_k$. Hence we have the following commutative diagram

$$\begin{array}{ccc} \mathbb{G}_k/B_k & \longrightarrow & U/B_k \cong BB_k \\ \downarrow & & \downarrow \\ Spec(k) \cong \mathbb{G}_k/G_k & \longrightarrow & U/G_k \cong BG_k. \end{array}$$

Since $CH^*(Spec(k))/p = 0$ for $* > 0$, we have the lemma. □

7. The orthogonal group $SO(m)$ and $p = 2$

We consider the orthogonal groups $G = SO(m)$ and $p = 2$. in this section. The mod 2-cohomology is written as (see for example [Mi-Tod], [Ni])

$$grH^*(SO(m); \mathbb{Z}/2) \cong \Lambda(x_1, x_2, ..., x_{m-1})$$

where $|x_i| = i$, and the multiplications are given by $x_s^2 = x_{2s}$. We write $y_{2(odd)} = x_{odd}^2$. Hence we can write

$$H^*(SO(m); \mathbb{Z}/2) \cong P(y) \otimes \Lambda(x_1, x_3, ...x_{\bar{m}}),$$

$$with \quad P(y) = \otimes_{i=0}^{s} \mathbb{Z}/2[y_{4i+2}]/(y_{4i+2}^{2^{r_i}}), \quad grP(y) \cong \Lambda(x_2, x_4, ...x_{m'})$$

for adequate integers $\bar{m}, m', s, r_i$. For ease of argument, we only consider in the case $m = 2\ell + 1$ so that

$$H^*(G; \mathbb{Z}/2) \cong P(y) \otimes \Lambda(x_1, x_3, ..., x_{2\ell-1})$$

$$grP(y)/2 \cong \Lambda(y_2, ..., y_{2\ell}), \quad letting\ y_{2i} = x_{2i} \ \ (hence\ y_{4i} = y_{2i}^2).$$

(Note that the suffix means its degree and it is used differently from other sections.)

The Steenrod operation is given as

$$Sq^k(x_i) = \binom{i}{k}(x_{i+k}).$$

The Q_i-operations are given by Nishimoto [Ni]

$$Q_n x_{2i-1} = y_{2i-2^{n+1}-2}, \qquad Q_n y_{2i} = 0.$$

Considering the map

$$U(\ell) \to SO(2\ell) \to SO(2\ell + 1),$$

we see that $b_i = c_i\ mod(2)$ for the transgression $d_{2i}(x_{2i-1}) = b_i$ and c_i which is the i-th elementary symmetric function on $S(t)$, from Proposition 6.1 in the preceding section. Moreover we see $Q_0(x_{2i-1}) = y_{2i}$ in $H^*(G; \mathbb{Z}/2)$. From Lemma 3.1 or Corollary 3.2, we have

$$2y_{2i} = c_i\ mod(4) \quad in\ H^*(G/T).$$

Indeed, the cohomology $H^*(G/T)_{(2)}$ is computed completely by Toda-Watanabe [To-Wa]

Theorem 7.1. *([To-Wa]) There are $y_{2i} \in H^*(G/T)$ for $1 \le i \le \ell$ such that $\pi^*(y_{2i}) = y_{2i}$ for $\pi : G \to G/T$, and that we have an isomorphism*

$$H^*(G/T) \cong \mathbb{Z}[t_i, y_{2i}]/(c_i - 2y_{2i}, J_{2i})$$

where $c_i = \sigma(t_1, ..., t_\ell)$ is the elementary symmetric polynomial, and (with $y_{2j} = 0$ for $j > \ell$)

$$J_{2i} = 1/4(\sum_{j=0}^{2i}(-1)^j c_j c_{2i-j}) = y_{4i} - \sum_{0<j<2i}(-1)^j y_{2j}y_{4i-2j}.$$

By using Nishimoto's result for Q_i-operation, we have

Corollary 7.2. *In $BP^*(G/T)/I_\infty^2$, the element*

$$b_i = 2y_{2i} + \sum v_n(y(2i + 2^{n+1} - 2))$$

is represented as an element in $BP^*(BT)$ where $\pi^* y(2s) = y_{2s}$ for the map $\pi : G \to G/T$, and $b_i = c_i$ in $BP^*(G/T)/I_\infty$.

It is known by Marlin and Merkurjev (see [To2] for details) that the torsion index of $SO(2\ell + 1)$ (and $SO(2\ell + 2)$) is 2^ℓ. Here we give an another proof.

Theorem 7.3. $t(G) = t(SO(2\ell + 1)) = 2^\ell$.

Proof. We consider in $H^*(G/T)$

$$b_1...b_\ell = c_1...c_\ell = (2y_2)(2y_4)...(2y_{2\ell}) = 2^\ell y_{top}$$

where $y_{top} = y_2...y_{2\ell}$. Hence $t(G) \le 2^\ell$.

Conversely, let $2^{\ell-1}y_{top} = t$ in $S(t)$. Then t in the ideal $(b_1, ..., b_\ell)$ in $S(t)$. So we can write $t = \sum b_i t(i)$. Then we have

$$2^{\ell-1}y_{top} = 2\sum y_{2i}t(i)$$

which implies $2^{\ell-2}y_{top} = \sum y_{2i}t(i)$ since $H^*(G/T)$ has no torsion. Continue this argument, we have a relation $y_{top} = \sum yt$ with $t \in S(t)$ where the number of y_{2s}in each monomial in y is less or equal to $\ell - 1$. This contradicts to $H^*(G/T)/2 \cong P(y) \otimes S(t)/(2, b)$. □

Recall that $W = W_{SO(2\ell+1)}(T)$ is the Weyl group. Then $W \cong S_k^\pm$ is generated by permutations and change of signs so that $|S_k^\pm| = 2^k k!$. Hence we have

$$H^*(BT)^W \cong \mathbb{Z}_{(2)}[p_1, ..., p_k] \subset H^*(BT) \cong \mathbb{Z}_{(2)}[t_1, ..., t_k],\ |t_i| = 2$$

where the Pontryagin class p_i is defined by $\Pi_i(1+t_i^2) = \sum_i p_i$. Consider the maps

$$\eta : T \overset{\eta_1}{\subset} U(\ell) \to SO(2\ell+1) \overset{\eta_2}{\to} U(2\ell+1).$$

Then $c_{2i}(\eta) = p_i \in CH^*(BT)^W$ which is the image from $c_{2i}(\eta_2) \in CH^*(BSO(2\ell+1))$.

On the other hand, $p_i = c_i(\eta_1)^2 \ mod(2)$, where $c_i(\eta_1) = \sigma_i$ is the elementary symmetric function in $S(t)$. From Lemma 6.5, the composition of the following maps

$$CH^*(BG_k)/2 \to CH^*(BB_k)/2 \to CH^*(X)/2$$

is zero for $* > 0$, we get $c_i(\eta_1)^2 = \sigma_i^2 = 0$ in $CH^*(X)/2$.

Theorem 7.4. *Let $(G,p) = (SO(2\ell+1), 2)$. Then there is an isomorphism*

$$CH^*(R(\mathbb{G}_k))/2 \cong \mathbb{Z}/2[c_1, ..., c_\ell]/(c_1^2, ..., c_\ell^2) = \Lambda(c_1, ..., c_\ell).$$

Proof. We only need to see that $c_1 ... c_\ell$ is non zero in $CH^*(X)/2$ from Lemma 5.5. Otherwise, it is represented by $2S(t)$ since $CH^*(X)$ is generated by elements from $S(t)$. It means that $2^{\ell-1} y_{top} = 1/2(c_1 ... c_\ell) \in S(t)$. Hence $t(G) < 2^\ell$ and a contradiction. □

Theorem 7.5. *Let $(G,p) = (SO(2\ell+1), 2)$ and $X = \mathbb{G}_k/B_k$. Then there is an isomorphism*

$$CH^*(X)/2 \cong S(t)/(2, c_1^2, ..., c_\ell^2).$$

Proof. We have the isomorphism

$$CH^*(X)/2 \cong CH^*(R(\mathbb{G}_k)) \otimes S(t)/(2, c_1, ..., c_\ell)$$

$$\cong \mathbb{Z}/2[c_1, ..., c_\ell]/(c_1^2, ..., c_\ell^2) \otimes S(t)/(c_1, ..., c_\ell) \cong gr(S(t)/(2, c_1^2, ..., c_\ell^2)).$$

Hence we have an additive isomorphism in this theorem.

Moreover we have a surjective ring map $S(t)/(2, c_1^2, ..., c_\ell^2) \twoheadrightarrow CH^*(X)/2$. From the additive isomorphism, its kernel is zero, which induces the ring isomorphism of the theorem. □

Corollary 7.6. *Let $(G', p) = (SO(2\ell), 2)$ and $X' = \mathbb{G}'_k/B_k$ so that $G' \subset G = SO(2\ell+1)$. Then $t(G') = 2^{\ell-1}$, and*

$$CH^*(R(\mathbb{G}'_k))/2 \cong CH^*(R(\mathbb{G}_k))/(2, c_\ell) \cong \Lambda(c_1, ..., c_{\ell-1}),$$

$$CH^*(X')/2 \cong CH^*(X)/(2, c_\ell) \cong S(t)/(2, c_1, ^2, ..., c_{\ell-1}^2, c_\ell).$$

Proof. This corollary is easily shown from $H^*(G'; \mathbb{Z}/2) \cong H^*(G; \mathbb{Z}/2)/(y_{2\ell})$. For example, $grP(y)' \cong \Lambda(y_2, ..., y_{2\ell-2})$ and $t(G') = 2^{\ell-1}$. □

From the above theorem and Proposition 6.2, the following corollary is immediate.

Corollary 7.7. *Let $G'' = Sp(2\ell+1)$ and $X'' = \mathbb{G}''_k/B_k$. Then the natural maps $G \to G'' \supset Sp(\ell)$ induce the isomorphisms*

$$CH^*(X)/2 \cong CH^*(X'')/(2, t_i | i > \ell) \cong H^*(Sp(\ell)/T; \mathbb{Z}/2) \cong H^*(Sp(\ell); \mathbb{Z}/2).$$

We can also prove Theorem 7.5 more directly by using $Sp(2\ell+1)$ instead of $U(2\ell+1)$.

We now study $CH^*(X|_K)/p$ for some interesting extension K over k. Let K be an extension of k such that X does not split over K but splits over an extension over K of degree ap, $(a,p)=1$. Suppose that

$$(*) \quad y_{2i} \in Res_K, \ for\ 1 \le i \le \ell - 1$$

where $Res_K = Im(res : CH^*(X|_K)/2 \to CH(\bar{X})/2)$. We want to consider in the case $y_{2\ell} \notin Res_K$.

Lemma 7.8. *Suppose $(*)$ and $\ell \neq 2^n - 1$ for $n > 0$. Then $y_{2\ell} \in Res_K$.*

Proof. We see that if $\ell \neq 2^n - 1$, then each $y_{2\ell}$ is a target of the Steenrod operation. Recall

$$Sq^{2k}(y_{2i}) = \binom{i}{k} y_{2(i+k)}.$$

It is well known that if $i = \sum i_s 2^s$ and $k = \sum k_s 2^s$ for $i_s, k_s = 0$ or 1, then (in $mod(2)$)

$$\binom{i}{k} = \binom{i_m}{k_m} ... \binom{i_s}{k_s} ... \binom{i_0}{k_0}.$$

Note that if $i = 2^n - 1$, then all $i_s = 1$ (for $s < n$). Otherwise there is s such that $i_s = 1$ but $i_{s-1} = 0$. Take $k = 2^{s-1}$ and $i' = i - 2^{s-1}$. Then $i' + k = i$ and

$$\binom{i'}{k} = \binom{i_m = 1}{0} ... \binom{i'_s = 0}{k_s = 0} \binom{1}{1} ... \binom{i_0}{0} = 1.$$

This means $Sq^{2k}(y_{2i'}) = y_{2i}$ if $i \neq 2^n - 1$. □

Lemma 7.9. *Suppose $(*)$ and $\ell = 2^n - 1$. Then all*

$$py_{2\ell},\ v_1 y_{2\ell},\ ...,\ v_{n-1} y_{2\ell} \quad in\ Im(res_\Omega)$$

where $res_\Omega : \Omega^*(X)/2 \to \Omega^*(\bar{X})/2$.

Proof. From Corollary 7.2, we see

$$b_{\ell-(2^j-1)} = 2y(2\ell - 2(2^j - 1)) + v_1 y(2\ell - 2(2^{j-1} - 1)) + ... + v_j(y(2\ell))$$
$$= v_j(y_{2\ell}) \quad mod(y_2, y_4, ..., y_{2\ell-2}).$$

Hence we have $res_\Omega(b_{\ell-(2^j-1)}) = v_j(y_{2\ell}) \ \ mod(y_2, y_4, ..., y_{2\ell-2})$. □

Thus we have

Theorem 7.10. *Suppose* $(*)$ *and* $\ell = 2^n - 1$. *Then*

$$CH^*(R(\mathbb{G}_k)|_K)/2 \cong \Lambda(y_2, ..., y_{2\ell-2}) \otimes CH^*(R_n)/2,$$

where $CH^*(R_n)/2 \cong \mathbb{Z}/2\{1, py, v_1y, ..., v_{n-1}y\}$. *Moreover we have*

$$Res_k^K(CH^*(R(\mathbb{G}_k))/2) \cong \ CH^*(R_n)/2 \subset CH^*(R(\mathbb{G}_k)|_K)/2.$$

Each generator maps as

$$b_j \mapsto \begin{cases} v_s y_\ell & if\ j = \ell - (p^s - 1) \\ 0 & otherwise. \end{cases}$$

At last of this section, we consider the case $\bar{X} = G/P$ with

$$G = SO(2\ell + 1) \quad and \quad P = U(\ell).$$

Let us write this X by Y, i.e. $Y = \mathbb{G}_k/P_k$. From the cofibering $SO(2\ell + 1) \to \bar{Y} \to BU(\ell)$, we have the spectral sequence

$$E_2^{*,*} \cong H^*(SO(2\ell + 1); \mathbb{Z}/2) \otimes H^*(BU(\ell))$$

$$\cong P(y) \otimes \Lambda(x_1, ..., x_{2\ell-1}) \otimes \mathbb{Z}/2[c_1, ..., c_\ell] \Longrightarrow H^*(\bar{Y}; \mathbb{Z}/2).$$

Here the differential is given as $d_{2i}(x_{2i-1}) = c_i$. Hence

$$H^*(\bar{Y}; \mathbb{Z}/2) \cong P(y).$$

This case is studied by Vishik [Vi] and Petrov [Pe] as maximal orthogonal (or quadratic) grassmannian. From Theorem 7.4, we have

Proposition 7.11. *Let* $G = Sin(2\ell+1)$ *and* $\mathbb{G}_k$ *be a versal* G_k*-torsor. Let* $Y = \mathbb{G}_k/U(\ell)_k$. *Then*

$$CH^*(Y)/2 \cong CH^*(R(\mathbb{G}_k))/2 \cong \Lambda(c_1, ..., c_\ell).$$

Remark. Petrov gets the similar and more general result [Pe], and this fact is pointed out by Nikita Karpenko to the author.

From above proposition, $CH^*(R(\mathbb{G}_k))/2$ has the ring structure in this case.

In [Vi], Vishik originally defined the J-invariant $J(q)$ of a quadratic form q which corresponds the quadratic grassmannian (see [Vi] for details) by

$$J(q) = \{i_k | y_{2i_k} \in Rec_{CH}\} \subset \{0, ..., \ell\}.$$

Let I be the fundamental ideal of the Witt ring $W(k)$ so that $grW(k) = \oplus_n I^n/I^{n+1} \cong K_*^M(k)/2$ where $K_*^M(k)$ is the Milnor K-theory of k. Smirnov and Vishik (Proposition 3.2.31 in [Sm-Vi]) proves that

$$q \in I^n \quad if\ and\ only\ if \quad \{0, ..., 2^{n-1} - 2\} \subset J(q).$$

Hence the condition $(*)$ in Theorem 7.10 is equivalent to $q \in I^n$ for the quadratic form q corresponding to $Y|_K$.

We also note that $G = Spin(m)$ cases corresponds to $q \in I^3$ from $1, y_2, y_4 \in Res_{CH}$ (see Lemma 8.1 below). This fact is of course, well known.

8. $Spin(m)$ AND $p = 2$

Throughout this section, let $p = 2$, $G = SO(2\ell + 1)$ and $G' = Spin(2\ell + 1)$. By definition, we have the 2 covering $\pi : G' \to G$. It is well known that $\pi^* : H^*(G/T) \cong H^*(G'/T')$ where T' is a maximal torus of G'. However the twisted flag varieties are not isomorphic. We will study these.

Let $2^t \le \ell < 2^{t+1}$, i.e. $t = [log_2\ell]$. The mod 2 cohomology is

$$H^*(G'; \mathbb{Z}/2) \cong H^*(G; \mathbb{Z}/2)/(x_1, y_1) \otimes \Lambda(z) \quad |z| = 2^{t+2} - 1$$

$$\cong P(y)' \otimes \Lambda(x_3, x_5, ..., x_{2\ell-1}) \otimes \Lambda(z)$$

where $P(y) \cong \mathbb{Z}/2[y_2]/(y_2^{2^{t+1}}) \otimes P(y)'$. (Here $d_{2^{t+2}}(z) = y_2^{2^{t+1}}$ for $0 \ne y \in H^2(B\mathbb{Z}/2; \mathbb{Z}/2)$ in the spectral sequence induced from the fibering $G' \to G \to B\mathbb{Z}/2$.) Hence

$$grP(y)' \cong \otimes_{2i \ne 2^j} \Lambda(y_{2i}) \cong \Lambda(y_6, y_{10}, y_{12}, ..., y_{2\ell}).$$

The Q_i operation for z is given by Nishimoto [Ni]

$$Q_0(z) = \sum_{i+j=2^{t+1}, i<j} y_{2i}y_{2j},$$

$$Q_n(z) = \sum_{i+j=2^{t+1}+2^{n+1}-2, i<j} y_{2i}y_{2j}, \quad for\ n \ge 1$$

We know that

$$grH^*(G/T)/2 \cong P(y)' \otimes \mathbb{Z}[y_2]/(y_2^{2^{t+1}}) \otimes S(t)/(2, c_1, c_2, ..., c_\ell)$$

$$grH^*(G'/T')/2 \cong P(y)' \otimes S(t')/(2, c_2, ..., c_\ell, c_1^{2^{t+1}}).$$

(Here $d_{2^{t+2}}(z) = c_1^{2^{t+1}}$ in the spectral sequence converging $H^*(G'/T')$.) These are additively isomorphic. Hence we have

Lemma 8.1. *The element $\pi^*(y_2) = c_1 \in S(t')$ and $\pi^*(t_j) = c_1 + t_j$ for $1 \le j \le \ell$.*

Take k such that $\mathbb{G}_k$ is a versal G_k-torsor so that $\mathbb{G}'_k$ is also a versal G'_k-torsor. Let us write $X = \mathbb{G}_k/B_k$ and $X' = \mathbb{G}'_k/B'_k$. Then

$$CH^*(\bar{R}(\mathbb{G}'_k)/2 \cong P(y)'/2, \quad and \quad CH^*(\bar{R}(\mathbb{G}_k))/2 \cong P(y)/2.$$

Theorem 8.2. *There are surjections*

$$\Lambda(c_2, ..., c_\ell) \twoheadrightarrow CH^*(R(\mathbb{G}'_k))/2 \twoheadrightarrow \mathbb{Z}/2\{1, c_2, ..., c_\ell\}.$$

Proof. From Lemma 5.3, we only need to show $b'_i \neq 0$ in $\Omega^*(G_k/T_k)/(I_\infty \cdot Im(res_\Omega))$. In fact, when $i \neq 2^j$, then in $H^*(G'/T')/4$, we have

$$2y_{2i} = c_j = b'_j \in S(t)$$

which is nonzero in $BP^*(G/T)/I_\infty \cdot Im(res_\Omega))$. Because $y_{2i} \in P(y)'$ and $y_{2i} \notin Im(res_{CH})$ from Lemma 4.5 since X is a versal flag variety.

When $i = 2^j$, we see $y_{2i} = y_{2^j} \in S(t')$, in fact $y_{2^j} \notin P(y)'$. Then in $BP^*(G'/T')/I_\infty^2$

$$2y_{2i}+v_1(y(2i+2))+...+v_n(y(2i+2^{n+1}-2)+... = c_i = b'_i \in BP^*(BT').$$

This element is nonzero in $BP^*(G/T)/I_\infty \cdot Im(res_\Omega))$ because

$$b'_i = v_1(y(2i+2)) \neq 0 \in k(1)^*(G/T)/(v_1 \cdot Im(res_{k(1)})$$

where $res_{k(1)} : k(1)^*(X') \to k(1)^*(\bar{X})$). Otherwise $y(2i+2) \in Im(res_{CH}))$, and this is a contradiction to $y_{2^j+2} \in Im(res_{CH})$, which follows from Lemma 4.5. □

Example. Let $G = SO(7)$ and $G' = Spin(7)$, i.e. $\ell = 3$. Their cohomologies are

$$H^*(G; \mathbb{Z}/2) \cong \mathbb{Z}/2[y_2, y_6]/(y_2^4, y_6^2) \otimes \Lambda(x_1, x_3, x_5),$$
$$H^*(G'; \mathbb{Z}/2) \cong \mathbb{Z}/2[y_6]/(y_6^2) \otimes \Lambda(x_3, x_5, z_7).$$

The cohomologies of flag manifolds are

$$H^*(G/T; \mathbb{Z}/2) \cong \mathbb{Z}/2[y_2, y_6]/(y_2^4, y_6^2) \otimes S(t)/(c_1, c_2, c_3),$$
$$H^*(G'/T'; \mathbb{Z}/2) \cong \mathbb{Z}/2[y_6]/(y_6^2) \otimes S(t)/(c_2, c_3, c_1^4).$$

These cohomologies are isomorphic by $\pi^*(y_2) = c_1$. The torsion indexes are $t(G) = 2^3$ and $t(G') = 2$. The Chow rings of versal flag varieties are

$$CH^*(X)/2 \cong S(t)/(2, c_1^2, c_2^2, c_3^2), \quad CH^*(R(\mathbb{G}_k))/2 \cong \Lambda(c_1, c_2, c_3),$$

$$CH^*(X')/2 \cong S(t)/(2, c_2^2, c_2c_3, c_3^2, c_1^4), \quad CH^*(R(\mathbb{G}'_k))/2 \cong \mathbb{Z}/2\{1, c_2, c_3\}.$$

Here $\pi^*(t_i) = c_1 - t_i$ so that $\pi^*(c_1) = 0 \ mod(2)$. For the third and the last isomorphisms, see Corollary 9.5 below. In fact G' is a group of type (I) and $CH^*(X')/(2, t_3) \cong CH^*(X'')/2$ where $X'' = \mathbb{G}_k/B_k$ for the exceptional Lie group G_2.

Marlin showed that

Lemma 8.3. *The torsion index $t(G')$ divides $2^{\ell-[log_2\ell]-1}$*

Proof. It follows from

$$\Pi_{i\neq 2^j} c_i = \Pi_{i\neq 2^j}(2y_{2i}) = 2^{\ell-t-1} y'_{top}$$

where y'_{top} is the generator of top degree elements in $P(y)'$. □

The exact value of $t(G')$ is determined Totaro, namely

$$t(G') = \ell - [log_2(\binom{\ell+1}{2} + 1)]$$

or that expression plus 1. (It is known $t(Spin(2\ell+1) = t(Spin(2\ell+2))$.

Marlin's bound fails first for $Spin(11)$. This fact was first founded by using a property of 12-dimensional quadrics [To2]. However we show it from the result from Nishimoto for Q_0-operation.

Lemma 8.4. *For $(G,p) = (Spin(11), 2)$, we have $t(G) = 2$ and*

$$CH^*(R(\mathbb{G}_k))/2 \cong \mathbb{Z}/2\{1, c_2, c_3, c_4, c_5, c_2c_4, c_1^8\}.$$

Proof. Recall the cohomology

$$H^*(G; \mathbb{Z}/2) \cong \mathbb{Z}/2[y_6, y_{10}]/(y_6^2, y_{10}^2) \otimes \Lambda(x_3, x_5, x_7, x_9, z_{15}).$$

By Nishimoto, we know $Q_0(z_{15}) = y_6y_{10}$. It implies $2y_6y_{10} = d_{16}(z_{15}) = c_1^8$, and hence $t(G) = 2$.

We can see $c_2c_3 = 0 \in CH^*(X)/2$. Because $c_2c_3 = 2v_1y_6^2 \in BP^*(G/T)$ and $2a \in Res_\Omega = Im(\Omega^*(X) \to BP^*(G/T)$ for $a \in BP^*(G/T)$, which implies $2y_6^2 \in Res_\Omega$. The cases $c_2c_5 = 0, c_2c_1^8 = 0, ...$ are proved similarly.

However note $c_2c_4 \neq 0 \in CH^*(X)/2$. Otherwise $v_1y_6y_{10}$ must be in Res_Ω. This means $v_1y_6y_{10} = b''$ for some $b'' \in BP^*(BT)$. However there is no $x \in H^{13}(G; \mathbb{Z}/2)$ such that $Q_1(x) = y_6y_{10}$ with $d_{12}(x) = b''$. □

In most cases, from the result Totaro, we see $\Pi_{i \neq 2^j} c_i = 0$. However from [To2] when $\ell = 8$, we know that $2^{\ell - [log_2(\ell)]-1} = 2^4 = t(Spin(17))$. (Note $y_{16} - 2y_6y_{10} \in S(t)$ but $y_{16} \notin S(t)$ when $\ell = 8$.) Hence we have

Corollary 8.5. *Let $\ell \geq 8$ and $G' = Spin(2\ell + 1)$. Let $X' = \mathbb{G}'_k/T'$. Then we have*

$$c_3c_5c_6c_7, \ c_3c_4c_6c_7 \neq 0 \in CH^*(X')/2.$$

Proof. It follows from that for $\ell = 8$, elements

$$c_3c_5c_6c_7 = 2^4y_6y_{10}y_{12}y_{14} = 2^4y'_{top} \quad and \quad c_3c_4c_6c_7 = 2^3v_1y'_{top}$$

are BP^*-module generators in $BP^*(G/T)/(I_\infty \cdot Im(res_\Omega))$. □

9. THE CASE E_8 AND $p = 5$

In this section, we consider the case $(G,p) = (E_8, 5)$. The similar arguments also hold for $(G,p) = (G_2, 2), (F_4, 3)$. The mod 5 cohomology of $G = E_8$ ([Mi-Tod]) is given by

Theorem 9.1. *The mod* 5 *cohomology* $H^*(E_8;\mathbb{Z}/5)$ *is isomorphic to*

$$\mathbb{Z}/5[y_{12}]/(y_{12}^5)\otimes\Lambda(z_3,z_{11},z_{15},z_{23},z_{27},z_{35},z_{39},z_{47})$$

where suffix means its degree. The cohomology operations are given

$$\beta(z_{11})=y_{12},\quad \beta(z_{23})=y_{12}^2,\quad \beta(z_{35})=y_{12}^3,\quad \beta(z_{47})=y_{12}^4,$$
$$P^1z_3=z_{11},\quad P^1z_{15}=z_{23},\quad P^1z_{27}=z_{35},\quad P^1z_{39}=z_{47}.$$

In §2, we use the notation such that $y_1=y_{12}$ and $x_1=z_3,...,x_8=z_{47}$. Hence we can rewrite the cohomology as

$$H^*(G;\mathbb{Z}/p)\cong\mathbb{Z}/p[y]/(y^p)\otimes\Lambda(x_1,...,x_{2p-2})$$

for $(G,p)=(E_8,5)$. The above isomorphism also holds for $(G,p)=(G_2,2),\ (F_4,3)$. So hereafter in this and next section, we assume (G,p) is one of $(G_2,2),\ (F_4,3)$ or $(E_8,5)$. The cohomology operations are given as

$$\beta: x_{2i}\mapsto y^i,\quad P^1: x_{2i-1}\mapsto x_{2i}\qquad for\ 1\le i\le p-1.$$

Hence the Q_i operations are given

$$Q_1(x_{2i-1})=Q_0(x_{2i})=y^i\quad for\ 1\le i\le p-1.$$

Therefore we have the following lemma, by using Lemma 3.1 or Corollary 3.2.

Lemma 9.2. *In* $BP^*(G/T)/I_\infty^2$, *we have*

$$py_i=b_{2i}\quad mod(b_2,b_4,...,b_{2i-2})\cdot S(t),$$
$$v_1y_i=b_{2i-1}\quad mod(b_1,b_2,...,b_{2i-3})\cdot S(t).$$

Proof. From Corollary 3.2, we see $py(2i)=b_{2i}\in BP^*(G/T)/I_\infty^2$

$$y(2i)=y^i+\sum yt\quad where\ y\in P(y),\ t\in S(t)\ |t|\ge 2.$$

By induction on i, we see $py^i\in(b_2,...b_{2i-2})\cdot S(t)$. Hence we get the first equation. The second equations follows similarly, from $v_1y(2i-1)=b_{2i-1}$ using Corollary 3.2. □

The fundamental class $f\in H^{2d}(G/T)$, $2d=dim(G/T)$ is represented $f=y^{p-1}t$ with $t=t_{top}\in S(t)$. Hence $pf\in S(t)$, which means $t(G)_{(p)}=p$.

Recall that $X=\mathbb{G}_k/B_k$ is the versal flag variety, and $\Omega^*(X)$ is the algebraic cobordism so that $\Omega^*(X)\otimes_{\Omega^*}\mathbb{Z}_{(p)}\cong CH^*(X)$. By Petrov-Semenov-Zainoulline, it is known when G is one of $(G_2,2),(F_4,3),(E_8,5)$, the motive $R(\mathbb{G}_k)$ in Theorem 4.3 is just the original Rost motive R_2 defined by Rost and Voevodsky. Recall Theorem 4.6. The restriction

$$res_{\Omega|R}:\Omega^*(R(\mathbb{G}_k))\to\Omega^*(\bar{R}(\mathbb{G}_k))$$

is injective. Hence the following restriction is also injective

$$res_\Omega : \Omega^*(X) \to \Omega^*(\bar{X}) \cong BP^*(G/T).$$

Corollary 9.3. *We see*

$$CH^*(R_2)/p \cong CH^*(R(\mathbb{G}_k)/p \cong \mathbb{Z}/p\{1, b_1, ..., b_{2p-2}.\}.$$

In particular, $b_s \neq 0 \in CH^*(X)/p)$. *Moreover for* $1 \leq s, r \leq 2p-2$, *we see* $b_s b_r = 0$ *in* $CH^*(X)/p$.

Proof. Recall Corollary 5.3. We prove $b_1 \neq 0 \in CH^*(X)$, and the other cases are proved similarly. Note $b_1 = v_1 y \in \Omega^*(\bar{X})$. If $b_1 \in BP^{<0} \cdot Im(res_\Omega)$, then $y \in Im(res_\Omega)$ and this is contradiction. So $b_1 \neq 0$ in

$$CH^*(X) \cong \Omega^*(X)/(BP^{<0} \cdot \Omega^*(X)) \cong Im(res_\Omega)/(BP^{<0} \cdot Im(res_\Omega)).$$

For the last isomorphism, we used the injectivity of res_Ω. We prove $b_1^2 = 0 \in CH^*(X)$. We see

$$b_1^2 = (v_1 y)^2 = v_1^2 y^2 = v_1 b_3 \in BP^*(G/T).$$

This element in $BP^{<0} \cdot Im(res_\Omega)$. Hence it is zero in $CH^*(X)$ as above. Other cases are proved similarly. □

Theorem 9.4. *([Ya4]) Let* $(G, p) = (G_2, 2)$. $(F_4, 3)$ *or* $(E_8, 5)$, *and let* $X = \mathbb{G}_k/T_k$. *Then there is an isomorphism*

$$CH^*(X)/p \cong S(t)/(p, b_i b_j | 1 \leq i, j \leq 2p-2).$$

Proof. From the preceding corollary we have the surjection

$$S(t)/(p, b_s b_r) \to CH^*(X)/p.$$

On the other hand, it is immediate that there is an additive isomorphism

$$S(t)/(p, b_i b_j) \cong \mathbb{Z}/p\{1, b_1, ..., b_{2p-2}\} \otimes S(t)/(p, b)$$

which injects to $\Omega^*(\bar{X})/(BP^{<0} \cdot Im(res_\Omega))$. Hence we have the theorem. □

Example. Let $G = F_4$ and $p = 3$. We note $G'' = Spin(9) \subset G$ and

$$H^*(BG'')/3 \cong H^*(BT'')^{W''}/3 \cong \mathbb{Z}/3[p_1, ..., p_4]$$

for the Pontryagin classes p_i. So $H^*(G''/T'')/3 \cong S(t)/(3, p_1, ..., p_4)$. By using the induced map from $G'' \subset G$, we can see $b_i = p_i$ in $CH^*(X)/3$. Hence

$$CH^*(X)/3 \cong S(t)/(3, p_i p_j | 0 \leq i, j \leq 4).$$

Moreover by Toda [Tod 1], it is known

$$Im(H^*(BG; \mathbb{Z}/3) \to H^*(BT; \mathbb{Z}/3)) \cong H^*(BT; \mathbb{Z}/3)^W$$

$$\cong \mathbb{Z}/3[p_1, p_2, \bar{p}_5, \bar{p}_9, \bar{p}_{12}]/(\bar{r}_{12})$$

for some $\bar{p}_i$ and $\bar{r}_{12}$, e.g. $\bar{p}_5 = p_3(p_2 - p_1^2) + p_4 p_1$. However it is known

$$p_1, p_2, \bar{p}_5 \notin Im(i^*_{CH} : CH^*(BG)/3 \to CH^*(BT)/3).$$

So we see that the subalgebra A of $S(t)/3$ generated by $p_i p_j$ for $0 \leq i, j \leq 4$ contains $Im(i^*_{CH})$. However note $\bar{p}_5 \in A$ but $\bar{p}_5 \notin Im(i^*_{CH})$.

Let G' be of type (I). Then it is well known ([Mi-Tod]) that there is a natural embedding $i : G \subset G'$ where $(G, p) = (G_2, 2), (F_4, 3)$ or $(E_8, 5)$ such that $i^* : H^*(G'; \mathbb{Z}/p) \to H^*(G; \mathbb{Z}/p)$ is surjetive. Moreover the polynomial parts $P(y)$ are isomorphic by this map i^*. This means $CH^*(\bar{R}(\mathbb{G}_k))) \cong CH^*(\bar{R}(\mathbb{G}'_k))$. This fact implies

$$CH^*(R(\mathbb{G}_k)) \cong CH^*(R(\mathbb{G}'_k))$$

by a theorem by Vishik and Zainoulline (Corollary 6 in [Vi-Za]). Thus we have

Corollary 9.5. *([Ya4]) Let G' be of type (I). Then there are isomorphisms*

$$CH^*(R(\mathbb{G}'_k))/p \cong \mathbb{Z}/p\{1, b_1, ..., b_{2p-2}\},$$
$$CH^*(X)/p \cong S(t)/(p, b_i b_j, b_k | 1 \leq i, j \leq 2p-2,\ 2p-1 \leq k \leq \ell).$$

Proof. Since $b_k = 0$ in $BP^*(G/T)/I_\infty \cong H^*(G/T)/p$, we can write

$$b_k = \sum p y^i t(i) + \sum v_1 y^i t(i)' \quad in\ BP^*(G/T)/I_\infty^2$$

where $t(i), t(i)' \in BP^* \otimes S(t)$. Take new b_k by $b_k - \sum b_{2i} t(i) - \sum b_{2i-1} t(i)'$. Then $b_k = 0$ in $BP^*(G/T)/I_\infty^2$. □

Example. Recall the case $(G', p) = (Spin(7), 2)$ and $(G, p) = (G_2, 2)$. Then we can take $b_1 = c_2$, $b_2 = c_3$, and $b_3 = c_1^4$, in fact

$$CH^*(X')/2 \cong S(t)/(c_2^2, c_2 c_3, c_3^2, c_1^4), \quad CH^*(X)/2 \cong CH^*(X')/(c_1).$$

10. THE CASE $G = E_8 \times E_8$ AND $p = 5$

In this section, we consider in the case $(G, p) = (E_8 \times E_8, p = 5)$. As the preceding section, we also consider in the cases $(G_2 \times G_2, p = 2)$ and $(F_4 \times F_4, p = 3)$.

Let $G = G_1 \times G_2$ for $G_1 \cong G_2$ such that (G_i, p) is one of $(G_2, 2), (F_4, 3)$ and $(E_8, 5)$. Hence

$$H^*(G_1 : \mathbb{Z}/p) \cong \mathbb{Z}/p[y]/(y^p) \otimes \Lambda(x_1, ...x_\ell) \quad \ell = 2p-2$$
$$H^*(G_2 : \mathbb{Z}/p) \cong \mathbb{Z}/p[y']/((y')^p) \otimes \Lambda(x'_1, ...x'_\ell) \quad \ell = 2p-2$$

By considering the spectral sequence induced from the fibering $G_1 \times G_2 \to G/T \to BT_1 \times BT_2$, we can compute

$$grH^*(G/T; \mathbb{Z}/p) \cong H^*(G_1/T_1; \mathbb{Z}/p) \otimes H^*(G_2/T_2 : \mathbb{Z}/p)$$

$$\cong \mathbb{Z}/p[y,y'] \otimes S(t) \otimes S(t')/(f, f', b_1, ..., b_\ell, b'_1, ..., b'_\ell).$$

Let $\mathbb{G}_k$ and $\mathbb{G}_{i,k}$ be a versal G_k-torsor and a versal $G_{i,k}$-torsor respectively. Let $X = \mathbb{G}_k/T_k$ and $X_i = \mathbb{G}_{i,k}/T_{i.k}$. Then the projection $G \to G_i$ induces the surjective maps

$$S(t) \otimes S(t') \to CH^*(X_1) \otimes CH^*(X_2) \to CH^*(X).$$

Lemma 10.1. *For $1 \le i, j \le p-1$, we have*

$$py^i(y')^j \notin res_{CH}(CH^*(X) \to CH^*(\bar{X})).$$

Proof. In the spectral sequence

$$E_2^{*,*} \cong H^*(BT; H^*(G_1 \times G_2; \mathbb{Z}/p)) \Longrightarrow H^*(G/T; \mathbb{Z}/p),$$

all nonzero differentials are $d_{|x_i|+1}(x_i) = b_i$ or $d_{|x'_i|+1}(x'_i) = b'_i$. Hence there is no $x \in H^*(G; \mathbb{Z}/p)$ such that $Q_0(x) = y^i(y')^j$ and $d_{|x|+1}(x) \neq 0 \in H^*(BT; \mathbb{Z}/p)$. From Lemma 3.1, we know $py^i(y')^j \notin H^*(BT)$. By Karpenko's theorem, $CH^*(X)$ is generated by elements in $H^*(BT) \cong S(t) \otimes S(t')$. Hence we have the lemma. □

Corollary 10.2. *For $1 \le i, j \le p-1$, the elements in $\Omega^*(\bar{X})$*

$$b_{2i}b'_{2i} = p^2 y^i(y')^j, \quad b_{2i-1}b'_{2j-1} = v_1^2 y^i(y')^j$$

$$b_{2i}b'_{2j-1} = b_{2i-1}b'_{2i} = pv_1 y^i(y')^j$$

are BP^*-module generators in $Im(res_\Omega(\Omega^*(X) \to \Omega^*(\bar{X}))$.

Proof. If $v_1 y^i(y')^j \in Im(res_\Omega)$, then applying r_{Δ_1}, we have $py^i(y')^j \in Im(res_\Omega)$, which contradicts to the preceding lemma. The other cases are proved similarly.. □

Theorem 10.3. *Let us write $B = S(t) \otimes S(t')/(p, b_s b_t, b'_s b'_t)$. Then there are surjective maps*

$$B \twoheadrightarrow CH^*(X)/p \twoheadrightarrow B/(b_{2i}b'_{2j-1} = b_{2i-1}b'_{2j})$$

where s, t, i, j range $1 \le s, t, 2i, 2j \le 2p-2$. We have also the surjective map

$$CH^*(R(\mathbb{G}_k))/p \twoheadrightarrow \mathbb{Z}/p\{1, b_s, b'_r, b_s b'_r\}/(b_{2i}b'_{2j-1} = b_{2i-1}b'_{2j}).$$

Proof. We can easily see that the right hand side ring in the theorem is isomorphic to the image

$$Im(res_\Omega)/I_\infty^2 \subset \Omega^*(\bar{X})/(I_\infty Im(res_k^{\bar{k}})).$$

□

Let K/k be a field extension of order p such that $\mathbb{G}_k$ is still non split over K, but for an extension K' over K of $order = ap$ with $(a,p)=1$, the group $\mathbb{G}_k$ is split. Consider the map for $K(X_1) \cong K(X_2)$

$$CH^*(X_1 \times X_2) \twoheadrightarrow CH^*(K(X_1) \times_K X_2) = CH^*(X_2|_{K(X_1)}) \cong CH^*(\bar{X}_2).$$

is surjective. In particular, there is

$$y'' \in CH^*((X_1 \times X_2)|_K)/p \quad such\ that \ \ y'' = y' \ in\ CH^*(\bar{X}_2)/p.$$

We consider the natural map $i : (G_1 \times G_1)/(T_1 \times T_1) \to G/T \times G/T$. Let us write $y_1 = i^*(y \otimes 1)$ and $y_2 = i^*(y')$. Then $J(\mathbb{G}_k) = (1,0)$ for $G = G_1 \times G_1$. Hence we have

Lemma 10.4. *Let $G = G_1 \times G_1$ for a type (I) group G_1. Let $X_1 = \mathbb{G}_{1,k}/T_k$ and $X_1 = (\mathbb{G}_1)_k/T_k$. Then we have*

$$CH^*(X|_K) \cong CH^*(X_1|_K) \otimes CH^*(\bar{X}_1),$$
$$CH^*(R(\mathbb{G}_k)|_K) \cong CH^*(R(\mathbb{G}_{1,k}) \otimes \mathbb{Z}/p[y'']/((y'')^p).$$

Theorem 10.5. *Let $G = G_1 \times ... \times G_m$ with $G_i \cong G_j$ for all i,j and $X = \mathbb{G}_k/T_k$. Let K/k be an extension of index pa with $(p,a) = 1$ and $CH^*(X|_K) \ncong CH^*(X)$. We have the nonzero torsion element $x(n) \in CH^*(X)$ such that in $CH^*(X)/p$*

$$x(n) = b_{2p-3}(1)....b_{2p-3}(n) \in Ker(res_k^K(CH^*(X)/p \to CH^*(X|_K)/p))$$

where $b_i(m) = pr_m^*(b_i)$ for the projection $pr_m : G \to G_m$.

Proof. The element $x(n)$ exists in $CH^*(X)$ since so does each $b_s(m)$. The element $x(n)$ is represented in $\Omega^*(\bar{X})$ by

$$v_1^n y(1)^{p-1}...y(n)^{p-1} = v_1^n y_{top} \quad for\ y(m) = pr_m^*(y).$$

But we see $x(n)/v_1$ is not in $res_k^{\bar{k}}(\Omega^*(X) \to \Omega(\bar{X}))$. Otherwise by using cohomology operation $r_{n\Delta_1}$, we see

$$p^{n-1} y_{top} \in Im(res_k^{\bar{k}}(CH^*(X)/p \to CH^*(\bar{X})/p).$$

The fundamental class of $\bar{X}$ is written as $f = y_{top}t$ for $t \in S(t)$. Hence above facts contradict to $t(G)_{(p)} = p^n$.

Thus $x(n)$ is an Ω^*-module generator in $res_k^{\bar{k}}(\Omega^*(X) \to \Omega^*(\bar{X}))$. This means $x(n) \neq 0 \in CH^*(X)/p$. (This element is torsion since it is zero in $CH^*(\bar{X}) \cong H^*(G/T)$.)

The fact $x(n)|_K = 0$ is proved similarly the preceding corollary. For example, take $y(n)'' = y(n)\ mod(y(i)|i < n)$ such that $y(n)'' \in Im(res_k^K)$ from the preceding lemma. Then

$$x(n)|_K = (b_{2p-3}(1)...b_{2p-3}(n))|_K = x(n-1)|_K \cdot v_1 y(n)''.$$

Hence it is zero in $\Omega^*(X|_K) \otimes_{\Omega^*} \mathbb{Z}/p \cong CH^*(X|_K)/p$. □

11. The case $G = E_8$ and $p = 3$

In this section, we assume $(G, p) = (E_8, p = 3)$. Recall the *mod* 3 cohomology of E_8 is given

Theorem 11.1. *The cohomology $H^*(E_8; \mathbb{Z}/3)$ is isomorphic to*

$$\mathbb{Z}/3[y_8, y_{20}]/(y_8^3, y_{20}^3) \otimes \Lambda(z_3, z_7, z_{15}, z_{19}, z_{27}, z_{35}, z_{39}, z_{47}).$$

Here the suffix means its degree, e.g., $|z_i| = i$.

By Kono-Mimura [Ko-Mi] the actions of cohomology operations are also known

Theorem 11.2. *([Ko-Mi]) We have $P^3 y_8 = y_{20}$, and*

$$\beta : z_7 \mapsto y_8, \quad z_{15} \mapsto y_8^2, \quad z_{19} \mapsto y_{20},$$

$$z_{27} \mapsto y_8 y_{20}, \quad z_{35} \mapsto y_8^2 y_{20}, \quad z_{39} \mapsto y_{20}^2, \quad z_{47} \mapsto y_8 y_{20}^2$$

$$P^1 : \ z_3 \mapsto z_7, \quad z_{15} \mapsto z_{19}, \quad z_{35} \mapsto z_{39}$$

$$P^3 : \ z_7 \mapsto z_{19}, \quad z_{15} \mapsto z_{27} \mapsto -z_{39}, \quad z_{35} \mapsto z_{47}.$$

We use notations $y = y_8, y' = y_{20}$, and $x_1 = z_3, ..., x_8 = z_{47}$. Then we can rewrite the isomorphism

$$grH^*(G/T)/3 \cong \mathbb{Z}/3[y, y']/(y^3, (y')^3) \otimes S(t)/(b_1, , ..., b_8).$$

From Lemma 3.4, we have

Corollary 11.3. *Element $v_1 y + v_2 y' = b_1 \in BP^*(G/T)/I_\infty^2$ is represented by an element in $BP^*(BT)$.*

For $k = i + 3j + 1$ with $0 \le i, j \le 2$ and $2 \le k \le 8$, we have $\beta(x_k) = y^i (y')^j$. Hence we have (from the proof of Lemma 5.2)

Corollary 11.4. *For all non-zero monomials $u \in P(y)/3$ except for $(yy')^2$, we see $3u \in S(t)$. That is, for $2 \le k = i+3j+1 \le 8$, $0 \le i \le 2$, we have*

$$b_k = b_{i+3j+1} = 3y_1^i (y')^j \quad in \ H^*(G/T)/(3^2).$$

Lemma 11.5. *Let $(G, p) = (E_8, 3)$ and $X = \mathbb{G}_k/T_k$. In $BP^*(X)$, there are $b_i \in S(t)$ such that $b_i \neq 0 \in CH^*(X)/3$ and in $BP^*(G/T)/I_\infty^2$*

$$b_k = b_{i+3j+1} = \begin{cases} v_1 y + v_2 y' & if \ k = 1 \\ 3y^i (y')^j & if \ 0 \le i \le 1, \ k \ge 2 \\ 3y^2 (y')^j + v_1 (y')^{j+1} & if \ i = 2. \end{cases}$$

Proof. Acting r_{Δ_1} on the equation $v_1y + v_2y' = b_1$ in $BP^*(X)/I_\infty^2$, we have

$$3y + v_1 r_{\Delta_1}(y) + v_2 r_{\Delta_1}(y') = r_{\Delta_1}(b_1).$$

Note $P^1(y_i) \in S(t)/3$ in $H^*(G/T; \mathbb{Z}/3)$ since they are primitive. Hence $v_i r_{\Delta_1}(y_i) \in BP^* \otimes S(t)\ mod(I_\infty^2)$. So we have $3y = b_2$ in $BP^*(G/T)/I_\infty^2$. Acting $r_{3\Delta_1}$ on the equation $3y = b_2 \in BP^*(X)/I_\infty^2$, we have $3y' = r_{3\Delta_1}(b_2)$, which is written by b_3.

Next we consider about $3y^2$ in $BP^*(X)/I_\infty^2$. Since $3y^2 = b_3$ in $H^*(X)/(9)$, we have

$$3y^2 + v_1(a_1) + v_2(a_2) = b_3 \quad in\ BP^*(X)/I_\infty^2.$$

Since $Q_1(x_3) = y'$, we can take $a_1 = y'$ by using the relation $v_1y + v_2y' = b_1$. (For example, when $a_1 = y' + yb$, we use $v_1yb = -v_2y'b$.) Since v_2a_2 is primitive in $k(2)^*(G/T)/(I_\infty^2)$ (Recall the proof of Lemma 3.4), we can take $a_2 = 0$. Otherwise if $a_2 = \sum y^i(y')^j b$, for $i = 1, 2$, then

$$v_2 y^i \otimes (y')^j b \neq 0 \in k(2)^*(G) \otimes k(2)^*(G/T).$$

Hence we get $3y^2 + v_1y' = b_3 \quad in\ BP^*(X)/I_\infty^2$.

Acting $r_{3\Delta_1}$ and $r_{6\Delta_1}$ on the above equation, we have the formulas for yy' and $(y')^2$. Here we used $r_{n\Delta_1}(y') \in BP^*(BT)/(I_\infty^2)$. since it is primitive.

Similar arguments work for the element y^2y', and we can see the formula for $y(y')^2$. □

Corollary 11.6. *The torsion index $t(E_8)_{(3)} = 3^2$.*

Proof. The fundamental class (localized at 3) is given as $y_{top}t = y^2(y')^2t$ for some $t \in S(t)$. Since $b_2b_8 = (3y)(3y(y')^2) = 3^2y_{top} \in S(t)$, we see $t(E_8)_{(3)} = 3$ or 3^2.

Suppose $t(E_8) = 3$, namely, $3y^2(y')^2 = b' \in S(t)$. From lemma 3.1, this implies that there is $x \in H^*(G; \mathbb{Z}/3)$ such that $Q_0(x) = y^2(y')^2$ and $d_r(x) = b'$. But such x does not exist from Theorem 12.2. □

Corollary 11.7. *Let $(G, p) = (E_8, 3)$ and $X = \mathbb{G}_k/T_k$ be a versal flag variety. Then b_1b_6, b_1b_8, b_2b_8 are non zero in $CH^*(X)/3$.*

Proof. First note that the fundamental class $f = (yy')^2t$ in $H^*(G/T)$ for some $t \in S(t)$. Since $t(E_8)_{(3)} = 3^2$ and X is a versal flag variety, we see $3(yy')^2 f \notin res_{CH}$. It follows $3(yy')^2 \notin res_{CH}$. Therefore $9(yy')^2, 3v_1(yy')^2, 3v_2(yy')^2$ are BP^*-module generators in Res_Ω, since

$$res_\Omega \ :\ (b_2b_8) \mapsto 9(yy')^2, \quad (b_1b_8) \mapsto 3v_1(yy')^2, \quad (b_1b_6) \mapsto 3v_2(yy')^2,$$

which shows they are non zero.

□

Note that for example, we can not see $b_1b_2 \neq 0$ from the above arguments. Indeed, $b_1b_2 = 3v_1y^2 = v_1(3y^2) = v_1b_3$ in $BP^*(G/T)/(I_\infty^2, v_2)$.

Giving a filtration on $S(t)$ by b_i, we have the isomorphism

$$grS(t)/3 \cong \mathbb{Z}/3[b_1, ..., b_8] \otimes S(t)/(b)$$

where $S(t)/(b) = S(t)/(3, b_1, ..., b_8)$. Recall that

$$A_N = \mathbb{Z}/3\{b_{i_1} \cdots b_{i_s} || b_{i_1}| + ... + |b_{i_s}| \leq N\}.$$

Since $CH^*(X) = 0$ if $2* > dim(G/T)$, we have the surjection

$$A_M \otimes S(t)/(b) \twoheadrightarrow CH^*(X)/3 \quad for\ M = |(yy')^2| = 56.$$

Theorem 11.8. *There are surjections*

$$A_{56} \twoheadrightarrow CH^*(R(\mathbb{G}_k))/3 \twoheadrightarrow \mathbb{Z}/3\{1, b_1, ..., b_8, b_1b_6, b_1b_8, b_2b_8\},$$

Proof. We get the result from the map

$$\Omega^*(R(\mathbb{G})) \to \Omega^*(\bar{R}(\mathbb{G}_k)) \cong \Omega^*[y, y']/(y^3, (y')^3)$$

and the preceding corollary. Indeed, elements $b_1^2, ..., b_2b_8$ are nonzero in $\Omega^*(\bar{R}(\mathbb{G}_k)/(3 \cdot Im(res_\Omega \Omega^*(R(\mathbb{G}_k))))$. □

We recall that there is an embedding $F_4 \subset E_8$ and $CH^*(\mathbb{G}'_k/B_k)/3 = CH^*(X')/3$ for $G' = F_4$ is given in Example in §9. Let K/k be a field extension of degree $3a$ with $(3, a) = 1$ such that the flag variety $X|_K = (\mathbb{G}_k/T_k)|_K$ is still twisted but $X|_{K'}$ is split for an extension K'/K of degree $3a'$ with $(3, a') = 1$. Note $P^3y = y'$ and if $y \in res_K^{\bar{K}}$, then so is y'. Since $X|_K$ is twisted, we see $y' \in res_K^{\bar{K}}$ but y is not. Hence the J-invariants are

$$J(\mathbb{G}_k|_K) = (1, 0) \quad but \quad J(\mathbb{G}_k) = (1, 1).$$

(See also 4.1.3 in [Pe-Se-Za], [Se] for E_8, $1 \geq j_1 \geq j_2$.)

We want to compute the Chow group of generalized Rost motive $R(\mathbb{G}_k|_K)$. Note that

$$CH^*(\bar{R}(\mathbb{G}_k|_K))/3 \cong \mathbb{Z}/3[y]/(y^3).$$

(while $CH^*(\bar{R}(\mathbb{G}_k))/3 \cong \mathbb{Z}/3[y, y']/(y^3, (y')^3)$) from th above J-invariants.

Let $R(\mathbb{F}_K)$ be the generalized Rost motive for the exceptional group $G = F_4$, Then it is known it is just the original Rost motive R_2 [Pe-Se-Za] and so $CH^*(\bar{R}(\mathbb{F}_K))/3 \cong \mathbb{Z}/3[y]/(y^3)$. Hence the natural map $i : F_4 \to E_8$ induces the isomorphism of motives over $\bar{K}$

$$CH^*(\bar{R}(\mathbb{F}_K))/3 \cong CH^*(\bar{R}(\mathbb{G}_k|_K)/3.$$

By a theorem by Vishik-Zainouline (Corollary 6 in [Vi-Za]), we have the isomorphism (without bar)

$$CH^*(R(\mathbb{F}_K))/3 \cong CH^*(R(\mathbb{G}_k|_K))/3.$$

It means that $R(\mathbb{G}_k|_K)$ is also the (original) Rost motive R_2. Thus we have

Lemma 11.9. *For $X = \mathbb{G}_k/B_k$, we have the additive isomorphism*

$$CH^*(X|_K))/3 \cong \mathbb{Z}/3\{1, b'_1, b'_2, b'_3, b'_4\} \otimes \mathbb{Z}/3[y']/((y')^3) \otimes S(t)/(b)$$

where $b'_1 = v_1 y, b'_2 = 3y, b'_3 = v_1 y^2, b'_4 = 3y^2$ in $BP^*(G/T)/I^2_\infty$.

Proposition 11.10. *Let us write the restriction map*

$$res^K_k : CH^*(R(\mathbb{G}_k))/3 \to CH^*(R(\mathbb{G}_k)|_K)/3$$
$$\cong CH^*(R(\mathbb{G}_k|_K)) \otimes \mathbb{Z}/3[y']/((y')^3).$$

Then we have $Im(res^K_k) \cong \mathbb{Z}/3\{1, b'_1, b'_2, b'_4, b'_2y', b'_4y', b'_2(y')^2\}$.

Proof. First note $b_4 = 3y', b_7 = 3(y')^2$ are in $Ker(res^K_k)$. Then we have

$$Im(res^K_k) \cong \mathbb{Z}/3\{1, b_1, b_2, b_3, b_5, b_6, b_8\}$$
$$\cong \mathbb{Z}/3\{1, v_1y, 3y, 3y^2, 3yy', 3y^2y', 3y(y')^2\}.$$

Here $b'_1 = v_1y, b'_2 = 3y, b'_4 = 3y^2, ..., b'_2(y')^2 = 3y(y')^2$ in $\Omega^*(X|_K)/I^2_\infty$. □

12. The case $G = E_8, p = 2$.

In this section, we consider the case $(G, p) = (E_8, 2)$. The *mod*(2) cohomology $H^*(E_8; \mathbb{Z}/2)$ is given as

$$\mathbb{Z}/2[z_3, z_5, z_9, x_{15}]/(z_3^{16}, z_5^8, z_9^4, z_{15}^4) \otimes \Lambda(z_{17}, z_{23}, z_{27}, z_{29}).$$

Here we consider a graded algebra $grH^*(E_8; Z/2)$ identifying $y_{2i} = z_i^2$ for $i = 3, 5, 9, 15$. Hereafter let $H^*(E_8; \mathbb{Z}/2)$ mean this $grH^*(E_8; \mathbb{Z}/2)$ as §7, 8.

Theorem 12.1. *The cohomology $H^*(E_8; \mathbb{Z}/2)$ is given*

$$\mathbb{Z}/2[y_6, y_{10}, y_{18}, y_{30}]/(y_6^8, y_{10}^4, y_{18}^2, y_{30}^2), \otimes \Lambda(z_3, z_5, z_9, z_{15}, z_{17}, z_{23}, z_{27}, z_{29}).$$

Let us write $y_1 = y_6, ..., y_4 = y_{30}$ and $x_1 = z_3, x_2 = z_5, ..., x_8 = z_{29}$. For ease of argument, let $x_4 = z_{17}$ and $x_5 = z_{15}$. Hence we can write

$$grH^*(E_8; \mathbb{Z}/2) \cong \mathbb{Z}/2[y_1, y_2, y_3, y_4]/(y_1^8, y_2^4, y_3^2, y_4^2) \otimes \Lambda(x_1, ..., x_8).$$

Lemma 12.2. *The cohomology operations acts as*

$$x_1 = z_3 \xrightarrow{Sq^2} x_2 = z_5 \xrightarrow{Sq^4} x_3 = z_9 \xrightarrow{Sq^8} x_4 = z_{17}$$
$$x_5 = z_{15} \xrightarrow{Sq^8} x_6 = z_{23} \xrightarrow{Sq^4} x_7 = z_{27} \xrightarrow{Sq^2} x_8 = z_{29}$$
$$x_5 = z_{15} \xrightarrow{Sq^2} x_4 = z_{17}$$

The Bockstein acts $Sq^1(x_{i+1}) = y_i$ for $1 \le i \le 3$, $Sq^1(x_8) = y_4$ and

$$Sq^1 \ : \ x_5 = z_{15} \mapsto y_1y_2, \quad x_6 = z_{23} \mapsto y_1y_3 + y_1^4, \quad x_7 = z_{27} \mapsto y_2y_3.$$

Then we see from Lemma 3.4

Corollary 12.3. *In $BP^*(X)/I_\infty^2$, we can take y_1 such that for $r_{2\Delta_1}(y_1) = y_2$ and $r_{4\Delta_1}(y_2) = y_3$, we have*

$$v_1y_1 + v_2y_2 + v_3y_3 = b_1 \quad for\ b_1 \in BP^*(BT).$$

From Lemma 3.1 and the Sq^1 action in Lemma 12.2, it is immediate that

Lemma 12.4. *Let $(G,p) = (E_8, 2)$ and $X = \mathbb{G}_k/T_k$. In $H^*(X)/(4)$, there are $b_i \in S(t)$ such that*

$$b_k = \begin{cases} 2y(1)\ (resp.\ 2y(2),\ 2y(3)) & if\ k = 2\ (resp.\ k = 3,4) \\ 2y(1,2)\ (resp.\ 2y(1,3),\ 2y(2,3)) & if\ k = 5\ (resp.\ k = 6,7) \\ 2y(4) & if\ k = 8. \end{cases}$$

where $y(i) = \pi^* y_i$, $y(i,j) = \pi^*(y_iy_j)$ for the map $\pi : G \to G/T$.

Lemma 12.5. *In $BP^*(X)/I_\infty^2$, we have*

$$b_i = \begin{cases} 2y_1 + v_2(y_1^2) + v_3(y_2^2) & if\ i = 2 \\ 2y_2 + v_1(y_1^2) + v_3(y_1^4) & if\ i = 3 \\ 2y_3 + v_1(y_2^2) & if\ i = 4 \end{cases}$$

Proof. On the equation $v_1y_1 + v_2y_2 + v_3y_3 = b$, we act the cohomology operation r_{Δ_1}, and we get

$$2y_1 + v_1 r_{\Delta_1}(y_1) + v_2 r_{\Delta_1}(y_2) + v_3 r_{\Delta_1}(y_3) = r_{\Delta_1}(b_1).$$

Recall that $P^1(y_i)$ are primitive in $H^*(G/T; \mathbb{Z}/2)$. In fact, by Kono-Ishitoya, we know (Theorem 5.9 in [Ko-Is])

$$P^1(y_1) \in S(t), \quad P^1(y_2) = y_1^2, \quad P^1(y_3) = y_2^2.$$

Thus we have $2y_1 + v_2(y_1^2) + v_3(y_2^2) = b_2$. (Note also $Q_2(x_2) = y_1^2$, $Q_3(x_2) = y_2^2$.)

Acting $r_{2\Delta_1}$ on this formula, we have

$$\begin{aligned} b_3 &= 2y_2 + v_1(y_1^2) + v_2(r_{\Delta_1}(y_1)^2) + v_3(r_{\Delta_1}(y_2)^2) \\ &= 2y_2 + v_1(y_1^2) + v_3(y_1^4). \end{aligned}$$

Acting $r_{4\Delta_1}$, we have $b_4 = 2y_3 + v_1y_2^2$ where we used $P^2(y_1) = y_2$. □

The anti-automorphism χ is defined by

$$\chi(Sq^0) = Sq^0, \quad \sum_i Sq^i \chi Sq^{n-i} = 0 \quad for\ n > 0.$$

For example, (when $Sq^1 = 0$) $\chi(P^i) = P^i$ for $i = 1, 2$ and

$$\chi(P^3) = P^3 + P^2P^1 + P^1P^2 = P^2P^1$$

since $P^3 = P^1P^2$, and $\chi(P^4) = P^4 + P^2P^2$.

From Lemma 3.1, we only see that

$$b = 2y(1,2) = 2(y_1y_2 + \lambda y_1^2 b') \quad where\ \lambda \in \mathbb{Z}/2,\ b, b' \in S(t).$$

However, it is known the more strong result

Lemma 12.6. *([Na], [To1]) In $H^*(G/T)/4$, we see $y_1y_2 \in S(t)$. Indeed, in the notation in [Na] (resp. [To1])*

$$\rho_8 = b + 2\gamma_3\gamma_5 \quad (resp.\ \ d_8 = 1/9d_4^2 - 2/3g_3g_5).$$

where $\gamma_3 = g_3 = y_1$, $\gamma_5 = g_5 = y_2$ and $d_i \in S(t)$.

Lemma 12.7. *In $BP^*(G/T)/(I_\infty^2)$, we have*

$$b_5 = 2y_1y_2 + v_1(y_3) + v_2(y_2b' + y_2^2b'') + v_3(y_4)$$

for some $b', b'' \in S(t)$.

Proof. From the preceding lemma, we can write

$$b_5 = 2y_1y_2 + v_1(a_1) + v_2(a_2) + v_3(a_3) \quad in\ BP^*(G/T)/(v_3, I_\infty^2).$$

We may assume that a_1 does not contains y_1 by using the relation b_1. Note that in $k(i)^*(G/T)/I_\infty^2$, each v_ia_i are primitive. Since y_2 is not in Q_1-image in $H^*(G;\mathbb{Z}/2)$, we see y_2 is v_1-torsion free. So if a_1 contains y_2, then v_1a_1 is not primitive in $k(1)^*(G)$, which is a contradiction. So a_1 contain only y_3, indeed $Q_1x_5 = y_3$ implies $a_1 = y_3$.

For a_2, we know that y_1, y_3 are not v_2-torsion. Therefore a_2 only contains y_2, that is,

$$a_2 = y_2b' + y_2^2b''\ mod(y_2^2) \quad for\ b', b'' \in S(t).$$

By the primitivity in $k(3)^*(G/T)$, the element a_3 only contains y_3, y_4. We know $Q_3(x_5) = y_4$. If $a_3 = v_1(y_4 + y_3b'')$, then let new y_4 be the element $y_4 + y_3b''$. Thus we have the result. □

Lemma 12.8. *In $BP^*(G/T)/(I_\infty^2, v_2, v_3)$, we have*

$$b_6 = 2(y_1y_3 + y_1^4 + y_1^2b''') \quad for\ b''' \in S(t)$$

Proof. We consider the action $r_{4\Delta_1}$ on b_5. First consider

$$r_{4\Delta_1}(y_1y_2) = r_{3\Delta_1}(y_1)r_{\Delta_1}(y_2) + r_{2\Delta_1}(y_1)r_{2\Delta_1}(y_2) + r_{\Delta_1}(y_1)r_{3\Delta}(y_2) + y_1r_{4\Delta}(y_2).$$

Here $r_{3\Delta_1} = P^2P^1\ mod(2)$. Hence we have with $mod(2)$

$$r_{3\Delta_1}(y_1)r_{\Delta_1}(y_2) = P^2P^1(y_1)P^1(y_2) = by_1^2.$$

$$r_{2\Delta_1}(y_1)r_{2\Delta_1}(y_2) = y_2b'', and\ \ 2y_2b'' \in S(t)$$

$$r_{\Delta_1}(y_1)r_{3\Delta}(y_2) = b''b''' \in S(t), \quad and \quad y_1r_{4\Delta_1}(y_2) = y_1y_3.$$

Hence $r_{4\Delta_1}(y_1y_2) = y_1y_3 + by_1^2\ mod(BP^* \otimes S(t))$.

Next consider

$$r_{4\Delta_1}(v_1y_3) = 2r_{3\Delta}(y_3) + v_1(r_{4\Delta_1}(y_3)).$$

Here with $mod(2)$ we see

$$r_{3\Delta}(y_3) = P^2P^1(y_3) = P^2(y_2^2) = y_1^4.$$

We also see $r_{4\Delta_1}(y_3) = P^4y_3 \in S(t)$ from the primitivity in $H^*(G/T;\mathbb{Z}/2)$.

At last we can see

$$r_{4\Delta_1}v_2(b'y_2 + b''y_2^2) = v_1r_{2\Delta_1}(b'y_2 + b''y_2^2) = 0 \quad mod(v_2).$$

Because if its contains v_1y_2 or $v_1y_2^2$, then it contradicts to the primitivity of $k(1)^*(G/T)$. If it contains v_1y, then it is in $Ideal(v_2)$ by the relation b_1. Thus we have the result ($with\ mo(v_2, v_3)$) of this lemma. □

Remark. Totaro (Lemma 4.4 in [To1] shows in $H^*(G/T)$

$$d_6^2 - 25/81d_4^3 + 2(15g_9g_3 + 1/3g_3^4 - 5/3g_5d_7 - 125/9g_5g_3d_4) + 2^2(-23/3g_3^2d_6) = 0$$

where $g_3 = y_1, g_5 = y_2, g_9 = y_3$ and $d_i \in S(t)$. Therefore we can take $b''' = 0$ in the preceding lemma. Totaro also gives explicit formula d_7, d_8 in $H^*(G/T)$. However we also give

Lemma 12.9. *In $H^*(G/T)/4$, we have $b_7 = 2(y_2y_3 + by_1^2 + b'y_2^2)$.*

Proof. We consider the action $r_{2\Delta_1}$ on b_6. Since b_6 contains no v_1a_1, there is no non-zero element in $BP^{<0}\cdot BP^*(G/T)$ such that $r_{2\Delta_1}(v_ia_i) \neq 0 \in H^*(G/T)/4$. Hence we only need to consider $r_{2\Delta_1}(y_1y_3 + y_1^4 + y_1^2b)$.

At first we compute

$$r_{2\Delta_1}(y_1y_3) = y_2y_3 + r_{\Delta_1}(y_1)r_{\Delta_1}(y_3) + y_1r_{2\Delta}(y_3) = y_2y_3 + b'y_2^2 \quad mod(S(t)).$$

We have $r_{2\Delta_1}(y_1^4) = 0$ by the Cartan formula. We see with $mod(2)$

$$r_{2\Delta_1}(by_1^2) = r_{2\Delta_1}(b)y_1^2 + b(r_{\Delta_1}(y_1)^2 = b'y_1^2 \quad modS(t).$$

□

Remark that by Totaro (Lemma 4.4 in [To1]), we know we can take $b = b' = 0$ in the above lemma. These facts follow from $P^1(y_1) = 0 \in H^*(G/T;\mathbb{Z}/2)$.

Now we study the torsion index. Recall

$$y_{top} = y_1^7y_2^3y_3y_4 \in P(y),$$

and t_{top} are generators of the top dimensional element in $S(t)/(b)$ so that $f = y_{top}t_{top}$ is the fundamental class of $H^{2d}(G/T)$.

Lemma 12.10. *(Totaro [To1]) We have $t(E_8)_{(2)} = 2^6$.*

Proof. We consider the element

$$\tilde{b} = b_5^3 b_6 b_4 b_8 = 2^6 (y_1 y_2)^3 (y_1 y_3 + y_1^4 + y_1^2 b'')(y_3)(y_4).$$

Here using $y_3^2 = b' \in S(t)\ mod(2)$, we have

$$(y_1 y_3 + y_1^4 + y_1^2 b'')y_3 = y_1^4 y_3 + (y_1 b' + y_1^2 b'')y_3.$$

Hence we can write

$$\tilde{b} = 2^6 (y_{top} + \sum yt) \quad for\ |t| > 0.$$

from Lemma 5.5, we see $t(E_8)_{(2)} \leq 2^6$.

Suppose $t(E_8)_{(2)} \leq 2^5$, that is, $2^5 f = 2^5 y_{top} t_{top} \in S(t)$. Then $2^5 f$ must be in ideal $I = (b_1, ..., b_8)$, and we can write for $b_i = 2y_{(i)}$

$$(*) \quad 2^5 f = \sum b_i t(i) = 2 \sum y_{(i)} t(i) \quad for\ t((i) \in S(t).$$

Since $H^*(G/T)$ has no torsion, we have $2^4 f = \sum y_{(i)} t(i)$.

Let us rewrite $s = \sum y_{(i)} t(i) = \sum_I y^I t(I)$ for a monomial $y^I = y_1^{i_1} ... y_4^{i_4} \in P(y)$ for $I = (i_1, ..., i_4)$, and $t(I) \in S(t)$. Then $s \in Ideal(2)$ implies each $t(I) \in Ideal(b_1, ..., b_8) \subset S(t)$, since $H^*(G/T)/2 \cong P(y) \otimes S(t)/(b)$. Continue this arguments, we have

$$(**) \quad f = \sum y^I t(I) \quad in\ H^*(G/T).$$

Consider this equation in $H^*(G/T)/2$, and we see $y_{top} = \sum y^I t(I)$, that is $y_{top} = y^I$.

To get $(**)$ from $(*)$, we exchanges b_i to $2y_{(i)}$ five times.

We note that the largest number of y_i's of monomials in $y_{(j)}$ is 1 or 2 except for $y_{(6)} = (y_1 y_3 + y_1^4 + y_1^2 b)$ where the number is 4. We easily see that $y_{(6)}$ appears as $y_{(i)}$ just one time in the process $(*)$ to $(**)$. We also see that $y_{(i)} = y_{(8)}$ just one time for the existence of y_4. Let us write by $\sharp_y(a)$ the number of y_i's in a. Then

$$\sharp_y(y_{(i_1)} ... y_{(i_5)}) \leq 2 \times 3 + 4 + 1 = 11.$$

On the other hand $\sharp_y(y_{top}) = 7 + 3 + 1 + 1 = 12$. Thus $t(E_8)_2 \geq 2^6$. $\square$

Lemma 12.11. *Let $(i_1, ..., i_k) \subset (4, 5, 5, 5, 6, 8)$. Then $\tilde{b} = b_{i_1} ... b_{i_k}$ is a non zero element in $CH^*(X)/2$.*

For example, $b_5^3 \neq 0 \in CH^*(X)/2$.

Let K be an extension of k such that X does not split over K but splits over an extension over K of degree ap, $(a, p) = 1$. Suppose that

$$(*) \quad y_1, y_2, y_3 \in Res_K, \quad but \quad y_4 \notin Res_K$$

where $Res_K = Im(res : CH^*(X|_K)/2 \to CH(\bar{X})/2)$. (Compare the condition $(*)$ in §7.) That is, $J(\mathbb{G}_k|_K) = (0,0,0,1)$ and such K exists (see [Pe-Se-Za], [Se]).

Lemma 12.12. *Suppose* $(*)$. *The elements* $2y_4$, v_1y_4, v_2y_4, v_3y_4 *in* $Im(res_k : \Omega^*(X)/I_\infty^2 \to BP^*(G/T)/I_\infty^2)$ *with modulo the* BP^**-subalgebra* M *generated by* y_1, y_2, y_3 *and* $S(t)$ *of* $BP^*(G/T)/I_\infty^2$.

Proof. From Corollary 3.2 or Lemma 12.7, with modulo M, we have

$$b_5 = v_3y_4, \quad b_6 = v_2y_4, \quad b_7 = v_1y_4, \quad b_8 = 2y_4.$$

□

Thus we have the following theorem by arguments similar to those to get Theorem 7.10. (The motive $R(\mathbb{G}_k)|_K$ in the theorem is an example of motives given in Lemma 8.4 in [Se].)

Theorem 12.13. *There is an isomorphism*

$$CH^*(R(\mathbb{G}_k)|_K)/2 \cong \mathbb{Z}/2[y_1, y_2, y_3]/(y_1^8, y_2^4, y_3^2) \otimes CH^*(R_4)/2,$$

where $CH^*(R_4)/2 \cong \mathbb{Z}/2\{1, 2y, v_1y, v_2y_2, v_3y\}$. *We have*

$$Res_k^K(CH^*(R(\mathbb{G}_k))/2) \cong CH^*(R_4)/2 \subset CH^*(R(\mathbb{G}_k)|_K)/2.$$

Each generator b_j *maps as*

$$b_j \mapsto \begin{cases} v_{8-j}y_4 & if\ 5 \le j \le 8 \\ 0 & if\ 1 \le j \le 4. \end{cases}$$

13. E_7 AND $p = 2$

The mod 2 cohomology of E_7 is given

$$H^*(E_7; \mathbb{Z}/2) \cong H^*(E_8)/(z_3^4, z_5^4, z_{15}^2, z_{29}).$$

We can write the notations in the preceding sections

Theorem 13.1. *We have an isomorphism*

$$H^*(E_7; \mathbb{Z}/2) \cong \mathbb{Z}/2[y_1, y_2, y_3]/(y_1^2, y_2^2, y_3^2) \otimes \Lambda(x_1, ..., x_7),$$

where $i^*(y_j) = y_j$ *for* $1 \le j \le 3$ *and* $i^*(x_i) = x_i$ *for* $1 \le i \le 7$ *and* $i^*(y_4) = i^*(x_8) = 0$ *for the natural embedding* $i : E_7 \subset E_8$

Corollary 13.2. *In* $BP^*(X)/I_\infty^2$, *we can take* y_1 *such that for* $r_{2\Delta_1}(y_1) = y_2$ *and* $r_{4\Delta_1}(y_2) = y_3$, *we have*

$$v_1y_1 + v_2y_2 + v_3y_3 = b_1 \quad for\ b_1 \in BP^*(BT).$$

From Lemma 3.1 and the Sq^1 action in Lemma 7.2, it is immediate

Lemma 13.3. *Let $(G,p)=(E_7,2)$. In $H^*(G/T)/(4)$, for all monomials $u\in P(y)/2$, except for $y_{top}=y_1y_2y_3$, the elements $2u$ are written as elements in $H^*(BT)$. Namely, in $H^*(G/T)/(4)$, there are $b_i\in S(t)$ such that*

$$b_k=\begin{cases}2y_1\ (resp.\ 2y_2,\ 2y_3) & if\ k=2\ (resp.\ k=3,4)\\ 2y_1y_2\ (resp.\ 2y_1y_3,\ 2y_2y_3) & if\ k=5\ (resp.\ k=6,7).\end{cases}$$

From lemma 10.5, it is immediate

Lemma 13.4. *In $BP^*(X)/I_\infty^2$, we have $2y_1=b_2, 2y_2=b_3, 2y_3=b_4$.*

Lemma 13.5. *We have $t(E_7)_{(2)}=2^2$.*

Proof. We get the result from $b_5b_3=(2y_1y_2)(2y_3)=2^2y_{top}$. □

Note that for $(G,p)=(E_7,2)$, the J-invarian $J(\mathbb{G}_k)=(1,1,1)$. Recall that $A_N=\mathbb{Z}/2\{b_{i_1}\cdots b_{i_s}||b_{i_1}|+...+|b_{i_s}|\le N\}$ and $|y_1y_2y_3|=34$.

Corollary 13.6. *There is a surjective maps*

$$A_{34}\twoheadrightarrow CH^*(R(\mathbb{G}_k))/2\twoheadrightarrow\mathbb{Z}/2\{1,b_1,...,b_7,b_1b_5,b_1b_6,b_1b_7\}.$$

Proof. In $\Omega^*(\bar{X})$, we see

$$b_1b_5=2v_3y_{top},\quad b_1b_6=2v_2y_{top},\quad b_1b_7=2v_1y_{top}.$$

These elements are Ω^*-module generators in $Im(res_k^{\bar{k}}(\Omega^*(X)\to\Omega(\bar{X}))$ because $2y_1y_2y_3\notin Im(res_k^{\bar{k}})$ from the fact $t(\mathbb{G}_k)=2^2$. □

Let us write $G'=E_8$ and $G''=G_2$ so that $G''\subset G=E_7\subset G'$. Take fields $k\subset K\subset K'$ such that

$$(**)\quad y_1^2,y_2^2,y_4\in Res_K\quad but\quad y_1,y_2,y_3\notin Res_K,$$

$$(***)\quad y_1^2,y_2,y_3,y_4\in Res_{K'}\quad but\quad y_1\notin Res_{K'}.$$

Then the following proposition is almost immediate

Proposition 13.7. *Suppose $(**)$ and $(***)$. We have isomorphisms,*

$$CH^*(R(\mathbb{G}'_k)|_K)/2\cong\mathbb{Z}/2[y_1^2,y_2^2,y_4]/(y_1^8,y_2^4,y_4^2)\otimes CH^*(\mathbb{G}_K)/2,$$

the restriction is given by $b_i\mapsto b_i$ for $1\le i\le 7$ and $b_8\mapsto 0$, and

$$CH^*(R(\mathbb{G}_K)|_{K'})/2\cong\mathbb{Z}/2[y_2,y_3]/(y_2^2,y_3^2)\otimes CH^*(R(\mathbb{G}'')|_{K'}))/2,$$

the restriction is given by $b_i\mapsto b_i$ for $i=1,2$, and $b_i\mapsto 0$ for $3\le i\le 7$.

References

[Ga-Me-Se] S. Garibaldi, A. Merkurjev and J-P Serre. Cohomological invariants in Galois cohomology. *University LecLet us writeture Series 28, Amer. math. Soc. Providence, RI* (2003), viii+168pp.

[Gr] A. Grothendieck. Torsion homologique et sections rationnelles. *Sem. C.Chevalley, ENS 1958,expose 5, Secreatariat Math., IHP, Paris* (1958).

[Ha] M.Hazewinkel, Formal groups and applications, *Pure and Applied Math. 78, Academic Press Inc.* (1978), xxii+573pp.

[Kar] N. Karpenko. Chow groups of some generically twisted flag variety. *www.ualberta.ca/"karpenko.* (2016).

[Ko-Is1] A. Kono and K. Ishitoya. Squaring operations in the 4-connective fiber spaces over the classifying spaces of the exceptional Lie groups. *RIMS Kyoto Univ. 91* (1985), 1299-1310.

[Ko-Is2] A. Kono and K. Ishitoya. Squaring operations in mod 2 cohomology of quotients of compact Lie groups by maximal tori. *Lect. Notes in Math. Springer 1298* (1987), 192-206.

[Ko-Mi] A. Kono and M. Mimura. Cohomology operations and the Hopf algebra structures of compact exceptional Lie groups E_7 and E_8. *Proc London Math. Soc.* **35** (1977), 345-358.

[Le-Mo1] M. Levine and F. Morel. Cobordisme algébrique I. *C. R. Acad. Sci. Paris* **332** (2001), 723-728.

[Le-Mo2] M. Levine and F. Morel. Cobordisme algébrique II. *C. R. Acad. Sci. Paris* **332** (2001), 815-820.

[Me-Ne-Za] A. Merkurjev, A. Neshitov and K. Zainoulline. *Invariants o degree* 3 *and torsion in the Chow group of a versal flag.* Composito Math. **151** (2015), 11416-1432.

[Me-Su] A. Merkurjev and A. Suslin. *Motivic cohomology of the simplicial motive of a Rost variety.* J. Pure and Appl. Algebra **214** (2010), 2017-2026.

[Mi-Ni] M. Mimura and T. Nishimoto. Hopf algebra structure of Morava K-theory of exceptional Lie groups. *Contem. Math.* **293** (2002), 195-231.

[Mi-Tod] M. Mimura and H. Toda, *Topology of Lie groups*, I and II, Translations of Math. Monographs, Amer. Math. Soc, **91** (1991).

[Na] M. Nakagawa. The integral cohomology ring of E_8/T. *Proc. Japan* (20), 383-394.

[Ni] T. Nishimoto. Higher torsion in Morava K-thoeory of $SO(m)$ and $Spin(m)$. *J.Math.Soc. Japan.* **52** (2001), 383-394.

[Pe] V. Petrov. Chow ring of generic maximal orthogonal grassmannians. *ftp://ftp.pdmi.ras.ru/pub/publicat/znsl/v443/p147.pdf* (2015).

[Pe-Se] V. Petrov, N. Semenov. Rost motives, affine varieties, and classifying spaces. *arXiv : 1506.007788v1.* (math. AT) Jun 2015.

[Pe-Se-Za] V.Petrov, N.Semenov and K.Zainoulline. J-Invariant of linear algebraic groups. *Ann. Scient. Ec. Norm. Sup.* **41**, (2008) 1023-1053.

[Ra] D.Ravenel. Complex cobordism and stable homotopy groups of spheres. *Pure and Applied Mathematics, 121. Academic Press* (1986).

[Ro1] M.Rost. Some new results on Chowgroups of quadrics. *preprint* (1990).

[Ro2] M.Rost. On the basic correspondence of a splitting variety. *preprint* (2006)

[Se] N. Semenov. Motivic construction of cohomological invariant. *Comment. Math. Helv. 91* (2016) 163-202.

[Se-Zh] N. Semenov and M Zhykhovich. Integral motives, relative Krull-Schumidt principle, and Maranda-type theorems. *Math. Ann. 363* (2015) 61-75.

[Sm-Vi] A. Smirnov and A. Vishik. Subtle characteristic classes. *arXiv : 1401.6661v1 [math.AG]* (2014).

[Su-Jo] A.Suslin and S.Joukhovistski. Norm Variety. *J.Pure and Appl. Algebra 206* (2006) 245-276.

[Tod1] H. Toda, Cohomology $mod(3)$ of the classifying space BF_4 of the exceptional group F_4. *J. Math. Kyoto Univ.* **13** (1973) 97-115..

[Tod2] H.Toda. On the cohomolgy ring of some homogeneous spaces. *J. Math. Kyoto Univ.* **15** (1975), 185-199.

[Tod-Wa] H.Toda and T.Watanabe. The integral cohomology ring of F_4/T and E_4/T. *J. Math. Kyoto Univ.* **14** (1974), 257-286.

[To1] B. Totaro. The torsion index of E_8 and the other groups. *Duke Math. J.* **299** (2005), 219-248.

[To2] B. Totaro. The torsion index of the spin groups. *Duke Math. J.* **299** (2005), 249-290.

[To3] B. Totaro, Group cohomology and algebraic cycles, *Cambridge tracts in Math. (Cambridge Univ. Press)* **204** (2014).

[Vi] A. Vishik. On the Chow groups of quadratic Grassmannians. *Doc. Math.* **10** (2005), 111-130.

[Vi-Ya] A.Vishik and N.Yagita. Algebraic cobordisms of a Pfister quadric. *London Math. Soc.* **76** (2007), 586-604.

[Vi-Za] A.Vishik and K.Zainoulline. Motivic splitting lemma. *Doc. Math.* **13** (2008), 81-96.

[Vo1] V. Voevodsky, The Milnor conjecture, *www.math.uiuc.edu/K-theory/0170* (1996).

[Vo2] V. Voevodsky. Motivic cohomology with $\mathbb{Z}/2$ coefficient. *Publ. Math. IHES* **98** (2003), 59-104.

[Vo3] V.Voevodsky. On motivic cohomology with $\mathbb{Z}/l$-coefficients. *Ann. of Math.* **174** (2011), 401-438.

[Ya1] N. Yagita. Algebraic cobordism of simply connected Lie groups. *Math.Proc.Camb.Phil.Soc.* **139** (2005), 243-260.

[Ya2] N. Yagita. Applications of Atiyah-Hirzebruch spectral sequence for motivic cobordism. *Proc. London Math. Soc.* **90** (2005) 783-816.

[Ya3] N. Yagita. Algebraic BP-theory and norm varieties. *Hokkaido Math. J.* **41** (2012), 275-316.

[Ya4] N. Yagita. Algebraic cobordism and flag varieties. *arXiv:1510.08135 (math. AT).* (2016)

FACULTY OF EDUCATION, IBARAKI UNIVERSITY, MITO, IBARAKI, JAPAN
E-mail address: nobuaki.yagita.math@vc.ibaraki.ac.jp,